

Stent recanalization of occluded iliofemoral and inferior cava veins in children with congenital heart disease: Acute and mid-term results.




Álvarez Fuente M., Zunzunegui J.L. Ballesteros F. Rodríguez Ogando A. Camino M. Maroto E.
Pediatric Cardiology, Hospital General Universitario Gregorio Marañón, Madrid, Spain

BACKGROUND:


Patients with congenital heart disease present a high risk of systemic vascular damage following catheterizations and indwelling lines. The limitation of vascular accesses can be problematic in these patients, requiring multiple surgeries and/or diagnostic and interventional catheterizations in the future. Our objectives are to evaluate the safety and efficacy of stenotic or occluded iliofemoral vein and inferior vena cava stent recanalization, and to determine the mid-term outcome following these procedures.

METHODS:

All patients with a vein occlusion and stent placed in the iliofemoral veins or in the inferior vena cava over a six year period (2004-2010) were included. Medical records and images were reviewed retrospectively in order to gather the following variables: age, weight, type and number of stents and follow-up procedures.



Right iliofemoral vein subtotal obstruction.



Everflex (6x40mm) stent implantation and distal balloon dilation.

RESULTS:

34 stents were placed in 17 patients (24 vessels).

Age: 5,7 years (1,3-10,1)

Weight: 19,2 Kg (4,4-34)


TOTAL UNILATERAL OCCLUSION	TOTAL BILATERAL OCCLUSION	SUBTOTAL UNILATERAL OCCLUSION	SUBTOTAL BILATERAL OCCLUSION	IVC
13	4	3	2	2

STENTS



Vessel Diameters:


Increase from 1,8 mm (1,1-2,5) to 8,8 mm (7,5-9,1).




There were no major procedure complications.

Patient follow-up:

- 4 patients (23%) underwent major surgery after recanalization.
- 9 patients (53%) had a total of 23 follow-up catheterizations, with a median length follow-up of 4 years.
- Two vessels (10,5%) were reoccluded requiring additional balloon dilation of existing stents.



Second Everflex (6x30mm) stent implantation.



Angiographic control.



36 month follow-up: recanalized iliofemoral vein. 8F catheter for endomyocardial biopsy.

CONCLUSIONS:

Stenotic or occluded vessel recanalization by intravascular stent placement is secure and effective to re-establish venous access for future catheterizations and/or surgeries. Reocclusion is unusual in our experience. Previously placed stents can be recanalized if necessary without major complications.