

Cardiovascular sequelae of Bronchopulmonary Dysplasia in Prematurely Born School Children with new echocardiography methods.

Bronchopulmonary dysplasia (BPD) is an important complication of prematurity and has long-term pulmonary consequences. Cardiovascular sequelae related to BPD have also been reported in severe BPD patients of the presurfactant era.

The aim of this study is to investigate cardiovascular consequences of BPD at school ages and to find out possible risk factors related to cardiovascular sequelae especially with new echocardiographic methods.

In our study totally 43 children (21F/22M: 23 aged 104 ± 7.4 months born preterm with BPD, 20 aged 112 ± 8.2 months born preterm without BPD and 21 healthy children aged 100 ± 16 months born at term (control group) were evaluated with conventional and myocardial tissue Doppler echocardiography. And also Left (LV) and Right ventricular (RV) strain and strain rate were assessed by speckle-tracking echocardiography.

Echocardiographic examination was performed using a Vivid 9 ultrasound machine, and LV and RV myocardial strain were analyzed offline using Echo PAC software.

BPD patients had a decreased pulmonary artery acceleration time and higher left and right ventricular myocardial performance indexes in school ages. Longitudinal and circumferential strains were abnormal in premature with BPD group even in the presence of normal RV and LVEF.

Negative effects of BPD on global cardiac performances of both ventricles and pulmonary arterial pressure persist up to school ages.