

**Institut für Phytopathologie
Christian-Albrechts-Universität Kiel**

**Bücher, Buchkapitel und begutachtete Publikationen
Books, book chapters and peer-reviewed publications**

2021

Begutachtete Publikationen:

1. **Birr, T., Hasler, M., Verreet, J.-A. & Klink, H. (2021):** Temporal changes in sensitivity of *Zymoseptoria tritici* field populations to different fungicidal modes of action. *Agriculture* 11, 269.
2. **Birr, T., Jensen, T., Preußke, N., Sönnichsen, F.D., De Boevre, M., De Saeger, S., Hasler, M., Verreet, J.-A. & Klink, H. (2021):** Occurrence of *Fusarium* mycotoxins and their modified forms in forage maize cultivars. *Toxins* 13, 110.
3. **Bonkowski, M., Tarkka, M., Razavi, B.S., Schmidt, H., Blagodatskaya, E., Koller, R., Yu, P., Knief, C., Hochholdinger, F. & Vetterlein, D. (2021):** Spatiotemporal dynamics of maize (*Zea mays* L.) root growth and its potential consequences for the assembly of the rhizosphere microbiota. *Frontiers in Microbiology* 12, 619499.
4. **de Mello, F.E., Lopes-Caitar, V.S., Prudente, H., Xavier-Valencio, S.A., Franzenburg, S., Mehl, A., Marcelino-Guimaraes, F.C., Verreet, J.-A., Balbi-Peña, M.I. & Godoy, C.V. (2021):** Sensitivity of *Cercospora* spp. from soybean to quinone outside inhibitors and methyl benzimidazole carbamate fungicides in Brazil. *Tropical Plant Pathology* 46, 69-80.
5. **Greenfield, L.M., Razavi, B.S., Bilyera, N., Zhang, X.C. & Jones, D.L. (2021):** Root hairs and protein addition to soil promote leucine aminopeptidase activity of *Hordeum vulgare* L. *Rhizosphere* 18, 100329.
6. **Guber, A., Blagodatskaya, E., Juyal, A., Razavi, B.S., Kuzyakov, Y. & Kravchenko, A. (2021):** Time-lapse approach to correct deficiencies of 2D soil zymography. *Soil Biology and Biochemistry* 157, 108225.
7. **Hashem, M.H. (2021):** Some scale insects and fungi infesting mango trees in Ismailia, Egypt. *Agricultura Tropica et Subtropica* 54, 136-146.
8. **Hosseini, S.S., Lakzian, A., Halajnia, A. & Razavi, B.S. (2021):** Optimization of EDTA and citric acid for risk assessment in the remediation of lead contaminated soil. *Rhizosphere* 17, 100277.
9. **Klink, H., Verreet, J.-A., Hasler, M. & Birr, T. (2021):** Will triazoles still be of importance in disease control of *Zymoseptoria tritici* in the future? *Agronomy* 11, 933.
10. **Liu, S., Zhang, S.Y., Dungait, J.A.J., Quine, T.A. & Razavi, B.S. (2021):** Rare microbial taxa rather than phoD gene abundance determine hotspots of alkaline phosphomonoesterase activity in the karst rhizosphere soil. *Biology and Fertility of Soils* 57, 257-268.
11. **Ma, Z.Z., Zhang, X.C., Zheng, B.Y., Yue, S.C., Zhang, X.C., Zhai, B.N., Wang, Z.H., Zheng, W., Li, Z.Y., Zamanian, K. & Razavi, B.S. (2021):** Effects of plastic and straw mulching on soil microbial P limitations in maize fields: Dependency on soil organic carbon demonstrated by ecoenzymatic stoichiometry. *Geoderma* 388, 114928.
12. **Meng, X.T., Liao, H.K., Fan, H.X., Zhang, X.C., Li, Y.Y., Yao, H.Y. & Razavi, B.S. (2021):** The geographical scale dependence of diazotroph assembly and activity: Effect of a decade fertilization. *Geoderma* 386, 114923.
13. **Pfeiffer, T., von Galen, A., Zink, P., Hübner, S., Linkies, A., Felgentreu, D., Drechsel, J., Birr, T., Röder, O., Kotte, M., Dietel, K., Junge, H., Schwarz, E.**

- & Koch, E. (2021):** Selection of bacteria and fungi for control of soilborne seedling diseases of maize. *Journal of Plant Diseases and Protection* 128, <https://doi.org/10.1007/s41348-021-00498-z>.
14. **Rezaei-Chiyaneh, E., Amirnia, R., Chiyaneh, S.F., Maggi, M., Barin, M. & Razavi, B.S. (2021):** Improvement of dragonhead (*Dracocephalum moldavica* L.) yield quality through coupled intercropping system and vermicompost application along with maintenance of soil microbial activity. *Land Degradation and Development* 32, 2833-2848.
 15. **Yu, P., He, X.M., Baer, M., Beirinckx, S., Tian, T., Moya, Y.A.T., Zhang, X.C., Deichmann, M., Frey, F.P., Bresgen, V., Li, C.J., Razavi, B.S., Schaaf, G., von Wirén, N., Su, Z., Bucher, M., Tsuda, K., Goormachtig, S., Chen, X.P. & Hochholdinger, F. (2021):** Plant flavones enrich rhizosphere bacteria to improve maize performance under nitrogen deprivation. *Nature Plants* 7, 481-499.
 16. **Zhang, X., Razavi, B.S., Liu, J., Wang, G., Zhang, X., Li, Z., Zhai, B., Wang, Z. & Zamanian, K (2021):** Croplands conversion to cash crops in dry regions: consequences of nitrogen losses and decreasing nitrogen use efficiency for food chain system. *Land Degradation and Development* 32, 1103-1113.
 17. **Zhang, X.C., Myrold, D.D., Shi, L.L., Kuzyakov, Y., Dai, H.C., Hoang, D.T.T., Dippold, M.A., Meng, X.T., Song, X.N., Li, Z.Y., Zhou, J. & Razavi, B.S. (2021):** Resistance of microbial community and its functional sensitivity in the rhizosphere hotspot to drought. *Soil Biology and Biochemistry* 161, 108360.

2020

Begutachtete Publikationen:

1. **Bilyera, N., Kuzyakova, I., Guber, A., Razavi, B.S. & Kuzyakov, Y. (2020):** How "hot" are hotspots: Statistically localizing the high-activity areas on soil and rhizosphere images. *Rhizosphere* 16, 100259.
2. **Birr, T., Hasler, M., Verreet, J.-A. & Klink, H. (2020):** Composition and predominance of *Fusarium* species causing Fusarium head blight in winter wheat grain depending on cultivar susceptibility and meteorological factors. *Microorganisms* 8, 617.
3. **Hamer, W.B., Birr, T., Verreet, J.-A., Duttmann, R. & Klink, H. (2020):** Spatio-temporal prediction of the epidemic spread of dangerous pathogens using machine learning methods. *ISPRS International Journal of Geo-Information* 9, 44.
4. **Hoang, D.T.T., Maranguit, D., Kuzyakov, Y. & Razavi, B.S. (2020):** Accelerated microbial activity, turnover and efficiency in the drilosphere is depth dependent. *Soil Biology and Biochemistry* 147, 107852, 107910.
5. **Jensen, T., de Boevre, M., de Saeger, S., Preußke, N., Sönnichsen, F.D., Kramer, E., Klink, H., Verreet, J.-A. & Birr, T. (2020):** Effect of ensiling duration on the fate of deoxynivalenol, zearalenone and their derivatives in maize silage. *Mycotoxin Research* 36, 127-136.
6. **Koch, E., Zink, P., Pfeiffer, T., von Galen, A., Linkies, A., Drechsel, J. & Birr, T. (2020):** Artificial inoculation methods for testing microorganisms as control agents of seed- and soil-borne Fusarium-seedling blight of maize. *Journal of Plant Diseases and Protection* 127, 883-893.
7. **Kravchenko, A.N., Guber, A.K., Razavi, B.S., Koestel, J., Quigley, M.Y., Robertson, G.P. & Kuzyakov, Y. (2020):** Reply to: "Variables in the effect of land use on soil extrapore enzymatic activity and carbon stabilization" by Glenn (2020). *Nature Communications*, 11, 6427.
8. **Lin, W.F., Zhang, H., Huang, D.M., Schenke, D., Cai, D.G., Wu, B.H. & Miao, Y. (2020):** Dual-localized WHIRLY1 affects salicylic acid biosynthesis via coordination of ISOCHORISMATE SYNTHASE1, PHENYLALANINE AMMONIA LYASE1 and S-

- ADENOSYL-L-METHIONINE-DEPENDENT METHYLTRANSFERASE1. *Plant Physiology* 184, 1884-1899.
9. **Liu, S.B., Pu, S.Y., Deng, D.L., Huang, H.Y., Yan, C., Ma, H. & Razavi, B.S. (2020):** Comparable effects of manure and its biochar on reducing soil Cr bioavailability and narrowing the rhizosphere extent of enzyme activities. *Environment International* 134, 105277.
 10. **Liu, S.B., Wang, J.Y., Pu, S.Y., Blagodatskaya, E., Kuzyakov, Y. & Razavi, B.S. (2020):** Impact of manure on soil biochemical properties: A global synthesis. *Science of the Total Environment* 745, 141003.
 11. **Ndossi, E.M., Becker, J.N., Hemp, A., Dippold, M.A., Kuzyakov, Y. & Razavi, B.S. (2020):** Effects of land use and elevation on the functional characteristics of soil enzymes at Mt. Kilimanjaro. *European Journal of Soil Biology* 97, 103167.
 12. **Pröbsting, M., Schenke, D., Hossain, R., Häder, C., Thurau, T., Wighardt, L., Schuster, A., Zhou, Z., Ye, W.Z., Rietz, S., Leckband, G. & Cai, D.G. (2020):** Loss-of-function of CRT1a (Calreticulin) reduces plant susceptibility to *Verticillium longisporum* in both *Arabidopsis thaliana* and oilseed rape (*Brassica napus*). *Plant Biotechnology Journal* 18, 2328-2344.
 13. **Schenke, D. (2020):** CRISPR/Cas9 or prime editing? - It depends on... *Open Life Sciences* 15, 868-870.
 14. **Schenke, D. & Cai, D.G. (2020):** Phytohormone crosstalk in the host-*Verticillium* interaction. *Plant Signaling and Behavior* 15, 1803567.
 15. **Schenke, D. & Cai, D.G. (2020):** Applications of CRISPR/Cas to improve crop disease resistance: Beyond inactivation of susceptibility factors. *iScience* 23, 101478.
 16. **Song, X.N., Razavi, B.S., Ludwig, B., Zamanian, K., Zang, H.D., Kuzyakov, Y., Dippold, M.A. & Gunina, A. (2020):** Combined biochar and nitrogen application stimulates enzyme activity and root plasticity. *Science of the Total Environment* 735, 139393.
 17. **Tian, P., Razavi, B.S., Zhang, X.C., Wang, Q.K. & Blagodatskaya, E. (2020):** Microbial growth and enzyme kinetics in rhizosphere hotspots are modulated by soil organics and nutrient availability. *Soil Biology and Biochemistry* 141, 107662.
 18. **Zhang, X., Kuzyakov, Y., Zang, H., Dippold, M.A., Shi, L., Spielvogel, S. & Razavi, B.S. (2020):** Rhizosphere hotspots: root hairs and warming control microbial efficiency, carbon utilization and energy production. *Soil Biology and Biochemistry* 148, 107872, 108061.

2019

Begutachtete Publikationen:

1. **Behrens, F.H., Schenke, D., Hossain, R., Ye, W.Z., Schemmel, M., Bergmann, T., Häder, C., Zhao, Y., Ladewig, L., Zhu, W.X. & Cai, D.G. (2019):** Suppression of abscisic acid biosynthesis at the early infection stage of *Verticillium longisporum* in oilseed rape (*Brassica napus*). *Molecular Plant Pathology* 20, 1645-1661.
2. **Birr, T., Verreet, J.-A. & Klink, H. (2019):** Prediction of deoxynivalenol and zearalenone in winter wheat grain in a maize-free crop rotation based on cultivar susceptibility and meteorological factors. *Journal of Plant Diseases and Protection* 126, 13-27.
3. **Duan, C.J., Razavi, B.S., Shen, G.T., Cui, Y.X., Yu, W.L., Li, S.Q. & Fang, L.C. (2019):** Deciphering the rhizobium inoculation effect on spatial distribution of phosphatase activity in the rhizosphere of alfalfa under copper stress. *Soil Biology and Biochemistry* 137, 107574.

4. **Jensen, T., de Boevre, M., Preußke, N., de Saeger, S., Birr, T., Verreet, J.-A. & Sönnichsen, F.D. (2019):** Evaluation of high-resolution mass spectrometry for the quantitative analysis of mycotoxins in complex feed matrices. *Toxins* 11, 531.
5. **Kravchenko, A.N., Guber, A.K., Razavi, B.S., Koestel, J., Quigley, M.Y., Robertson, G.P. & Kuzyakov, Y. (2019):** Microbial spatial footprint as a driver of soil carbon stabilization. *Nature Communications* 10, 3121.
6. **Kuzyakov, Y. & Razavi, B.S. (2019):** Rhizosphere size and shape: Temporal dynamics and spatial stationarity. *Soil Biology and Biochemistry* 135, 343-360.
7. **Razavi, B.S., Zhang, X.C., Bilyera, N., Guber, A. & Zarebanadkouki, M. (2019):** Soil zymography: Simple and reliable? Review of current knowledge and optimization of the method. *Rhizosphere* 11, 10016.
8. **Schenke, D., Utami, H.P., Zhou, Z., Gallegos, M.-T. & Cai, D.G. (2019):** Suppression of UV-B stress induced flavonoids by biotic stress: Is there reciprocal crosstalk? *Plant Physiology and Biochemistry* 134, 53-63.
9. **Wei, L., Razavi, B.S., Wang, W.Q., Zhu, Z.K., Liu, S.L., Wu, J.S., Kuzyakov, Y. & Ge, T.D. (2019):** Labile carbon matters more than temperature for enzyme activity in paddy soil. *Soil Biology and Biochemistry* 135, 134-143.
10. **Yuan, H.Z., Liu, S.L., Razavi, B.S., Zhran, M., Wang, J.R., Zhu, Z.K., Wu, J.S. & Ge, T.D. (2019):** Differentiated response of plant and microbial C: N: P stoichiometries to phosphorus application in phosphorus-limited paddy soil. *European Journal of Soil Biology* 95, 103122.
11. **Zhong, X.B., Zhou, Q.Z., Cui, N., Cai, D.G. & Tang, G.X. (2019):** BvcZR3 and BvHs1pro-1 genes pyramiding enhanced beet cyst nematode (*Heterodera schachtii* Schm.) resistance in oilseed rape (*Brassica napus* L.). *International Journal of Molecular Sciences* 20, 1740.

2018

Begutachtete Publikationen:

1. **Conrad, N., Brandes, M., Will, T., Verreet, J.-A., Ulber, B. & Heimbach, U. (2018):** Effects of insecticidal seed treatments and foliar sprays in winter oilseed rape in autumn on insect pests and TuYV infection. *Journal of Plant Diseases and Protection* 125, 557-565.
2. **Rudelt, J., Klink, H. & Verreet, J.-A. (2018):** Einfluss der Zugabe adhäsiver Additive in Beizlösungen auf das Fließverhalten von Getreidesaatgut. *Journal für Kulturpflanzen* 70, 158-162.
3. **Seychelles, L.H., Happe, S., Palacios, E., Ludwig, M., Hollmer, S., Ehlers, R.-U., Schulz, C. & Mercier, L. (2018):** Successful rearing of whiteleg shrimp *Litopenaeus vannamei* larvae fed a desiccation-tolerant nematode to replace *Artemia*. *Aquaculture Nutrition* 24, 903-910.
4. **Thomsen, G., Birr, T., Klink, H. & Verreet, J.-A. (2018):** Geographic distribution and disease severity of different *Rhizoctonia* species pathogenic towards maize in different regions in Germany and France. *Journal of Plant Diseases and Protection* 125, 461-468.
5. **Tillessen, A., Menkhaus, J. & Verreet, J.-A. (2017):** Development of specific PCR primers for diagnosis and quantitative detection of the fungal maize pathogen *Kabatiella zaeae*. *European Journal of Plant Pathology* 152, 503-506.

2017

Begutachtete Publikationen:

1. **Guo, L.B., Qiu, J., Ye, C.Y., Jin, G.L., Mao, L.F., Zhang, H.Q., Yang, X.F., Peng, Q., Wang, Y.Y., Jia, L., Lin, Z.X., Li, G.M., Fu, F., Liu, C., Chen, L., Shen, E.H.,**

Wang W.D., Chu, Q.J., Wu, D.Y., Wu, S.L., Xia, C.Y., Zhang, Y.F., Zhou, X.M., Wang, L.F., Wu, L.M., Song, W.J., Wang, Y.F., Shu, Q.Y., Aoki, D., Yumoto, E., Yokota, T., Miyamoto, K., Okada, K., Kim, D.-S., Cai, D.G., Zhang, C.L., Lou, Y.G., Qian, Q., Yamaguchi, H., Yamane, H., Kong, C.-H., Timko, M.P., Bai, L.Y. & Fan, L.J. (2017): *Echinochloa crus-galli* genome analysis provides insight into its adaptation and invasiveness as a weed. *Nature Communications* 8, 1031.

2. **Rudelt, J., Klink, H. & Verreet, J.-A. (2017):** Einfluss der Zusammensetzung der Beizlösung auf die Staubentwicklung an Getreidesaatgut. *Journal für Kulturpflanzen* 69, 303-308.
3. **Shen, Y.F., Sun, S., Hua, S.J., Shen, E.H., Ye, C.-Y., Cai, D.G., Timko, M.P., Zhu, Q.-H. & Fan, L.J. (2017):** Analysis of transcriptional and epigenetic changes in hybrid vigor of allopolyploid *Brassica napus* uncovers key roles for small RNAs. *Plant Journal* 91, 874-893.
4. **Ullmann, I., Herrmann, A., Hasler, M. & Taube, F. (2017):** Influence of the critical phase of stem elongation on yield and forage quality of perennial ryegrass genotypes in the first reproductive growth. *Field Crops Research* 205, 23-33.
5. **Zhou, Z., Schenke, D., Miao, Y. & Cai, D.G. (2017):** Investigation of the crosstalk between the flg22 and the UV-B-induced flavonol pathway in *Arabidopsis thaliana* seedlings. *Plant, Cell and Environment* 40, 453-458.

2016

Begutachtete Publikationen:

1. **Hamer, W.B., Verreet, J.-A. & Duttmann, R. (2016):** Räumliche und zeitliche Vorhersage der Eintrittswahrscheinlichkeit eines ertragsgefährdenden Mehltauereignisses an Winterweizen mit der Random-Forest-Methode. *Journal für Angewandte Geoinformatik* 2, 342-352.
2. **Hamer, W.B., Verreet, J.-A. & Duttmann, R. (2016):** Spatial prediction of the infestation risk of winter wheat by the pathogen *Blumeria graminis* f. sp. *tritici* (powdery mildew) in Schleswig-Holstein using semi-empirical and machine learning techniques. *GIS Science* 29, 140-148.
3. **Rüstner, W.K., Klink, H. & Verreet, J.-A. (2016):** Nachweis einer ALS-resistenten *Lolium perenne*-Population in Norddeutschland. *Journal für Kulturpflanzen* 68, 117-124.
4. **Ullmann, I., Herrmann, A., Hasler, M., Cai, D.G. & Taube, F. (2016):** Variability in the critical phase of stem elongation of perennial ryegrass genotypes: a source for breeding progress? *Journal of Agricultural Science* 154, 1002-1014.

2015

Bücher/Buchkapitel:

1. **Verreet, J.-A. (2015):** Die Agrar- und Ernährungswissenschaftliche Fakultät. In: O. Auge (ed.): *Christian-Albrechts-Universität zu Kiel – 350 Jahre Wirken in Stadt, Land und Welt*, 313-328. Wachholtz, Kiel, Hamburg.

Begutachtete Publikationen:

1. **Guo, L.B., Qiu, J., Han, Z.J., Ye, Z.H., Chen, C., Liu, C.J., Xin, X.F., Ye, C.-Y., Wang, Y.-Y., Xie, H.Q., Wang, Y., Bao, J.D., Tang, S., Xu, J., Gui, Y.J., Fu, F., Wang, W.D., Zhang, X.C., Zhu, Q.H., Guang, X.M., Wang, C.Z., Cui, H.F., Cai, D.G., Ge, S., Tuskan, G.A., Yang, X.H., Qian, Q., He, S.Y., Wang, J., Zhou, X.-P. & Fan, L.J. (2015):** A host plant genome (*Zizania latifolia*) after a century-long endophyte infection. *Plant Journal* 83, 600-609.

2. **Shen, E.H., Zou, J., Behrens, F.H., Chen, L., Ye, C.Y., Dai, S.T., Li, R.Y., Ni, M., Jiang, X.X., Qiu, J., Liu, Y., Wang, W.D., Zhu, Q.-H., Chalhoub, B., Bancroft, I., Meng, J.L., Cai, D.G. & Fan, L.J. (2015):** Identification, evolution, and expression partitioning of miRNAs in allopolyploid *Brassica napus*. *Journal of Experimental Botany* 66, 7241-7253.
3. **Tillner, R., Assheuer, T., Rennert, B., Trubiroha, A., Clemmesen, C. & Wuertz, S. (2015):** Evaluation of an improved RNA/DNA quantification method in a common carp (*Cyprinus carpio* Linnaeus 1758) larval feeding trial with *Artemia*, two nematodes (*Panagrellus redivivus* Linnaeus 1758, *Panagrolaimus* sp. Fuchs 1930) and dry feed. *Journal of Ichthyology* 31, 466-473.

2014

Begutachtete Publikationen:

1. **Effendi, Y., Radatz, K., Labusch, C., Rietz, S., Wimalasekera, R., Helizon, H., Zeidler, M. & Scherer, G.F.E. (2014):** Mutants of phospholipase A (pPLA-I) have a red light and auxin phenotype. *Plant Cell and Environment* 37, 1626-1640.
2. **Schenke, D. & Cai, D.G. (2014):** The interplay of transcription factors in suppression of UV-B induced flavonol accumulation by flg22. *Plant Signaling and Behavior* 9, e28745.
3. **Schenke, D., Cai, D.G. & Scheel, D. (2014):** Suppression of UV-B stress responses by flg22 is regulated at the chromatin level via histone modification. *Plant, Cell and Environment* 37, 1716-1721.
4. **Shen, D., Suhrkamp, I., Wang, Y., Liu, S., Menkhaus, J., Verreet, J.-A., Fan, L. & Cai, D.G. (2014):** Identification and characterization of microRNAs in oilseed rape (*Brassica napus*) responsive to infection with the pathogenic fungus *Verticillium longisporum* using *Brassica* AA (*Brassica rapa*) and CC (*Brassica oleracea*) as reference genomes. *New Phytologist* 204, 577-594.
5. **Shen, D. & Wu, B. (2014):** Structure, biological properties and utilities of marine-derived antimicrobial peptides. *Current Organic Chemistry* 18, 793-803.

2013

Bücher/Buchkapitel:

1. **Poehling, J.-M. & Verreet, J.-A., eds. (2013):** Lehrbuch der Phytomedizin, 4. Auflage. Eugen Ulmer, Stuttgart.

Begutachtete Publikationen:

1. **Anbesse, S. & Ehlers, R.-U. (2013):** *Heterorhabditis* sp. not attracted to synthetic (*E*)- β -caryophyllene, a volatile emitted by roots upon feeding by corn rootworm. *Journal of Applied Entomology* 137, 88-96.
2. **Anbesse, S., Sumaya, N.H., Dörfler, A.V., Strauch, O. & Ehlers, R.-U. (2013):** Stabilisation of heat tolerance traits in *Heterorhabditis bacteriophora* through selective breeding and creation of inbred lines in liquid culture. *Biocontrol* 58, 85-93.
3. **Anbesse, S., Sumaya, N.H., Dörfler, A.V., Strauch, O. & Ehlers, R.-U. (2013):** Selective breeding for desiccation tolerance in liquid culture provides genetically stable inbred lines of the entomopathogenic nematode *Heterorhabditis bacteriophora*. *Applied Microbiology and Biotechnology* 97, 731-739.
4. **Ayub, F., Strauch, O., Seychelles, L. & Ehlers, R.-U. (2013):** Influence of temperature on life history traits of the free-living, bacterial-feeding nematode *Panagrolaimus* sp. strain NFS-24. *Nematology* 15, 939-946.
5. **Dong, S.Q., Tian, Z.H., Chen, P.J., Kumar, R.S., Shen, C.H., Cai, D.G., Oelmüller, R. & Yeh, K.W. (2013):** The maturation zone is an important target of *Piriformospora indica* in Chinese cabbage roots. *Journal of Experimental Botany* 64,

- 4529-4540.
6. **Honnens, H., Assheuer, T. & Ehlers, R.-U. (2013):** Desiccation and storage of *Panagrolaimus* sp. (strain NFS-24-5). *Nematology* 15, 557-566.
 7. **Honnens, H. & Ehlers, R.-U. (2013):** Liquid culture of *Panagrolaimus* sp. for use as food for marine aquaculture shrimp and fish species. *Nematology* 15, 417-429.
 8. **Honnens, H. & Ehlers, R.-U. (2013):** Cultivation of a free-living nematode *Panagrolaimus* sp. in batch and fed-batch liquid culture of *Saccharomyces cerevisiae* for larval food in marine aquaculture. *Russian Journal of Nematology* 21, 13-21.
 9. **Nimkingrat, P., Khanam, S., Strauch, O. & Ehlers, R.-U. (2013):** Hybridisation and selective breeding for improvement of low temperature activity of the entomopathogenic nematode *Steinernema feltiae*. *Biocontrol* 58, 417-426.
 10. **Nimkingrat, P., Strauch, O. & Ehlers, R.-U. (2013):** Hybridisation and genetic selection for improving desiccation tolerance of the entomopathogenic nematode *Steinernema feltiae*. *Biocontrol Science and Technology* 23, 348-361.
 11. **Nimkingrat, P., Uhlmann, F., Strauch, O. & Ehlers, R.-U. (2013):** Desiccation tolerance of dauers of entomopathogenic nematodes of the genus *Steinernema*. *Nematology* 15, 451-458.
 12. **Seychelles, L., Doiron, K., Audet, C., Tremblay, R., Pernet, F. & Lemarchand, K. (2013):** Impact of arachidonic acid enrichment of live rotifer prey on bacterial communities in rotifer and larval fish cultures. *Canadian Journal of Microbiology* 59, 189-196.
 13. **Thurau, T., Beyer, M., Blanck, T. & Liu, X. (2013):** Transcriptional changes of putative *Fusarium graminearum* transporter sequences in response to trifloxystrobin and deoxynivalenol. *Journal of Plant Pathology* 95, S1.29-S1.37.
 14. **Zinkernagel, V. & Verreet, J.-A. (2013):** Günter Martin Hoffmann Obituary. *Journal of Plant Diseases and Protection* 120, 95-96.
 15. **Zinkernagel, V. & Verreet, J.-A. (2013):** Günter Martin Hoffmann – ein Nachruf. *Journal für Kulturpflanzen* 65, 334-335.

2012

Bücher/Buchkapitel:

1. **Verreet, J.-A., Klink, H., Birr, T. & Urban, K. (2012):** Fusarium-Pilze an Mais. In: BASF SE (ed.): Mais – Potenziale einer bedeutenden Kultur richtig nutzen!, 150-165. Ludwigshafen, BASF SE.
2. **Verreet, J.-A. & Knott, J. (2012):** Das Risiko von Fusarium in Mais – Mykotoxine im Erntegut. In: BASF SE (ed.): Mais – Potenziale einer bedeutenden Kultur richtig nutzen!, 166-171. Ludwigshafen, BASF SE.

Begutachtete Publikationen:

1. **Anbesse, S.A., Strauch, O. & Ehlers, R.-U. (2012):** Genetic improvement of the biological control nematode *Heterorhabditis bacteriophora* (Rhabditidomorpha: Heterorhabditidae): heterosis effect enhances desiccation but not heat tolerance through selective breeding and creation of inbred lines in liquid culture. *Biocontrol Science and Technology* 22, 1035-1045.
2. **Bakhali, A.H., Abd-Elsalam, K.A., Guo, J.R., Khiyami, M.A. & Verreet, J.-A. (2012):** Characterization of novel di-, tri-, and tetranucleotide microsatellite primers suitable for genotyping various plant pathogenic fungi with special emphasis on fusaria and *Mycosphaerella graminicola*. *International Journal of Molecular Sciences* 132, 2951-2964.
3. **Bao, J.D., Yao, J.Q., Zhu, J.Q., Hu, W.M., Cai, D.G., Li, Y., Shu, Q.Y. & Fan, L.J. (2012):** Identification of glutinous maize landraces and inbred lines with altered transcription of waxy gene. *Molecular Breeding* 30, 1707-1714.

4. **Hammoudi, O., Salman, M., Abuamsha, R. & Ehlers, R.-U. (2012):** Effectiveness of bacterial and fungal isolates to control *Phoma lingam* on oilseed rape *Brassica napus*. American Journal of Plant Sciences 3, 773-779.
5. **Kumar, A., Bhandari, A., Sinha, R., Sardar, P., Sushma, Goyal, P., Goswami, C. & Grapputo, A. (2012):** Molecular phylogeny of OVOL genes illustrates a conserved C2H2 zinc finger domain coupled by hypervariable unstructured regions. PLOS One 7, e39399.
6. **Rietz, S., Bernsdorff, F.E.M. & Cai, D.G. (2012):** Members of the germin-like protein family in *Brassica napus* are candidates for the initiation of an oxidative burst that impedes pathogenesis of *Sclerotinia sclerotiorum*. Journal of Experimental Botany 63, 5507-5519.
7. **Sardar, P., Kumar, A., Bhandari, A. & Goswami, C. (2012):** Conservation of tubulin-binding sequences in TRPV1 throughout evolution. PLOS One 7, e31488.
8. **Tamiru, T., Waeyenberge, L., Hailu, T., Ehlers, R.-U., Puza, V. & Mracek, Z. (2012):** *Steinernema ethiopiense* sp. n. (Rhabditida: Steinernematidae), a new entomopathogenic nematode from Ethiopia. Nematology 14, 741-757.
9. **Wilson, M.J., Ehlers, R.-U. & Glazer, I. (2012):** Entomopathogenic nematode foraging strategies – is *Steinernema carpocapsae* really an ambush forager? Nematology 14, 389-394.

2011

Bücher/Buchkapitel:

1. **Ehlers, R.-U., ed. (2011):** Regulation of Biological Control Agents. Springer, Dordrecht, Heidelberg, London, New York.
2. **Ehlers, R.-U. (2011):** Regulation of biological control agents and the EU policy support action REBECA. In: R.-U. Ehlers (ed.): Regulation of Biological Control Agents, 3-23. Springer, Dordrecht, Heidelberg, London, New York.
3. **Ehlers, R.-U. (2011):** Cost-benefit, risk and trade-off analysis of regulation. In: R.-U. Ehlers (ed.): Regulation of Biological Control Agents, 139-153. Springer, Dordrecht, Heidelberg, London, New York.
4. **Fjelsted, A. & Ehlers, R.-U. (2011):** Proposals on how to accelerate registration of biological control agents. In: R.-U. Ehlers (ed.): Regulation of Biological Control Agents, 375-405. Springer, Dordrecht, Heidelberg, London, New York.
5. **Strauch, O., Strasser, H., Hauschild, R. & Ehlers, R.-U. (2011):** Proposals for bacterial and fungal biocontrol agents. In: R.-U. Ehlers (ed.): Regulation of Biological Control Agents, 267-288. Springer, Dordrecht, Heidelberg, London, New York.

Begutachtete Publikationen:

1. **Abd-El Salam, K.A., Bakhali, A.H., Moslem, M., DeWit, P.J.G.M. & Verreet, J.-A. (2011):** Detection of *Mycosphaerella graminicola* in wheat leaves by a microsatellite dinucleotide specific-primer. International Journal of Molecular Sciences 12, 682-693.
2. **Abd-El Salam, K.A., Bakhali, A.H., Moslem, M. & Verreet, J.-A. (2011):** Development of di- and tetranucleotide repeat primer for discrimination of *Fusarium* species. Journal of Food Safety 31, 35-40.
3. **Abuamsha, R., Salman, M. & Ehlers, R.-U. (2011):** Improvement of seed bio-priming of oilseed rape (*Brassica napus* ssp. *oleifera*) with *Serratia plymuthica* and *Pseudomonas chlororaphis*. Biocontrol Science and Technology 21, 199-213.
4. **Abuamsha, R., Salman, M. & Ehlers, R.-U. (2011):** Effect of seed priming with *Serratia plymuthica* and *Pseudomonas chlororaphis* to control *Leptosphaeria maculans* in different oilseed rape cultivars. European Journal of Plant Pathology 130, 287-295.

5. **Abuamsha, R., Salman, M. & Ehlers, R.-U. (2011):** Differential resistance of oilseed rape cultivars (*Brassica napus* ssp. *oleifera*) to *Verticillium longisporum* infection is affected by rhizosphere colonisation with antagonistic bacteria, *Serratia plymuthica* and *Pseudomonas chlororaphis*. *Biocontrol* 56, 101-112.
6. **Addis, T., Mulawarman, M., Waeyenberge, L., Moens, M., Viaene, N. & Ehlers, R.-U. (2011):** Identification and intraspecific variability of *Steinernema feltiae* strains from Cemoro Lawang village in Eastern Java, Indonesia. *Russian Journal of Nematology* 19, 21-29.
7. **Beyer, M., Kiesner, F., Verreet, J.-A. & Klink, H. (2011):** Fungicide sensitivity of *Septoria tritici* field isolates is affected by an interaction between fungicidal mode of action and time. *Journal of Plant Pathology* 93, S1.7-S1.13.
8. **Fan, L.J., Bo, S.P., Chen, H.A., Ye, W.Z., Kleinschmidt, K., Baumann, H.I., Imhoff, J.F., Kleine, M. & Cai, D.G. (2011):** Genome sequence of *Bacillus subtilis* subsp. *spizizenii* gtP20b isolated from the Indian Ocean. *Journal of Bacteriology* 193, 1276-1277.
9. **Fan, L.J., Gui, Y.J., Zheng, Y.F., Wang, Y., Cai, D.G. & Yu, X.L. (2011):** Ancient DNA sequences of rice from the low Yangtze reveal significant genotypic divergence. *Chinese Science Bulletin* 56, 28-29.
10. **Fan, L.J., Yun, L., Li, Z.F., Baumann, H.I., Kleinschmidt, K., Ye, W.Z., Imhoff, J.F., Kleine, M. & Cai, D.G. (2011):** Draft genome sequence of the marine *Streptomyces* sp. strain PP-C42, isolated from the Baltic sea. *Journal of Bacteriology* 193, 3691-3692.
11. **Lee, Y.-C., Johnson, J.M., Chien, C.-T., Sun, S., Cai, D.G., Lou, B.G., Oelmüller, R. & Yeh, K.-W. (2011):** Growth promotion of Chinese cabbage and *Arabidopsis* by *Piriformospora indica* is not stimulated by mycelium-synthesized auxin. *Molecular Plant-Microbe Interactions* 24, 421-431.
12. **Mei, J., Qian, L., Disi, J.O., Yang, X., Li, Q., Li, J., Frauen, M., Cai, D.G. & Qian, W. (2011):** Identification of resistant sources against *Sclerotinia sclerotiorum* in *Brassica* species with emphasis on *B. oleracea*. *Euphytica* 177, 393-399.
13. **Nimkingrat, P., Ehlers, R.-U. & Strauch, O. (2011):** Desiccation tolerance among different isolates of the entomopathogenic nematode *Steinernema feltiae* (Filipjev). *Communications in Agricultural and Applied Biological Sciences* 76 (3), 293-296.
14. **Susurluk, I.A. (2011):** Potential of *Steinernema feltiae* (Rhabditida: Steinernematidae) as a biological control agent against the cabbage maggot *Delia radicum* (Diptera: Anthomyiidae) in oilseed rape. *Turkish Journal of Agriculture and Forestry* 35, 413-419.
15. **Tang, G., Knecht, K., Yang, X., Qin, Y., Zhou, W. & Cai, D.G. (2011):** A two-step protocol for shoot regeneration from hypocotyl explants of oilseed rape and its application for *Agrobacterium*-mediated transformation. *Biologia Plantarum* 55, 21-26.

2010

Bücher/Buchkapitel:

1. **Ehlers, R.-U. (2010):** REBECA – EU-policy support action to review regulation of biological control agents. In: U. Gisi, I. Chet, M.L. Gullino (eds.): *Recent Developments in Management of Plant Diseases - Plant Pathology in the 21st Century*, 147-161. Springer, Dordrecht, Heidelberg, London, New York.
2. **Thurau, T., Ye, W.Z. & Cai, D.G. (2010):** Insect and nematode resistance. In: F. Kempken, C. Jung (eds.): *Genetic Modification of Plants - Agriculture, Horticulture and Forestry*, 177-198. Springer, Berlin, Heidelberg, New York.

Begutachtete Publikationen:

1. **Chan, Y.-L., Cai, D.G., Taylor, P.W.J., Chan, M.-T. & Yeh, K.W. (2010):** Adverse effect of the chitinolytic enzyme PjCHI-1 in transgenic tomato on egg mass production and embryonic development of *Meloidogyne incognita*. *Plant Pathology* 59, 922-930.
2. **Hirao, A. & Ehlers, R.-U. (2010):** Influence of inoculum density on population dynamics and dauer juvenile yields in liquid culture of biocontrol nematodes *Steinernema carpocapsae* and *S. feltiae* (Nematoda: Rhabditida). *Applied Microbiology and Biotechnology* 85, 507-515.
3. **Hirao, A., Ehlers, R.-U. & Strauch, O. (2010):** Life cycle and population development of the entomopathogenic nematodes *Steinernema carpocapsae* and *S. feltiae* (Nematoda, Rhabditida) in monoxenic liquid culture. *Nematology* 12, 201-210.
4. **Knecht, K., Seyffarth, M., Desel, C., Thurau, T., Sherameti, I., Lou, B., Oelmüller, R. & Cai, D.G. (2010):** Expression of *BvGLP-1* Encoding a germin-like protein from sugar beet in *Arabidopsis thaliana* leads to resistance against phytopathogenic fungi. *Molecular Plant-Microbe Interactions* 23, 446-457.
5. **Kontradowitz, L. & Verreet, J.-A. (2010):** Assessment of resistance and virulence in the pathosystem sugar beet (*Beta vulgaris*)/powdery mildew (*Erysiphe betae*) - development of basics for an effective powdery mildew resistance breeding. *Journal of Plant Diseases and Protection* 117, 49-54.
6. **Mukuka, J., Strauch, O., Al Zainab, M.H. & Ehlers, R.-U. (2010):** Effect of temperature and desiccation stress on infectivity of stress tolerant hybrid strains of *Heterorhabditis bacteriophora*. *Russian Journal of Nematology* 18, 111-116.
7. **Mukuka, J., Strauch, O. & Ehlers, R.-U. (2010):** Variability in desiccation tolerance among different strains of the entomopathogenic nematode *Heterorhabditis bacteriophora*. *Nematology* 12, 711-720.
8. **Mukuka, J., Strauch, O., Hoppe, C. & Ehlers, R.-U. (2010):** Improvement of heat and desiccation tolerance in *Heterorhabditis bacteriophora* through cross-breeding of tolerant strains and successive genetic selection. *Biocontrol* 55, 511-521.
9. **Mukuka, J., Strauch, O., Hoppe, C. & Ehlers, R.-U. (2010):** Fitness of heat and desiccation tolerant hybrid strains of *Heterorhabditis bacteriophora* (Rhabditidomorpha: Heterorhabditidae). *Journal of Pest Science* 83, 281-287.
10. **Mukuka, J., Strauch, O., Waeyenberge, L., Viaene, N., Moens, M. & Ehlers, R.U. (2010):** Heat tolerance among different strains of the entomopathogenic nematode *Heterorhabditis bacteriophora*. *Biocontrol* 55, 423-434.
11. **Navaneethan, T., Strauch, O., Besse, S., Bonhomme, A. & Ehlers, R.-U. (2010):** Influence of humidity and a surfactant-polymer-formulation on the control potential of the entomopathogenic nematode *Steinernema feltiae* against diapausing codling moth larvae (*Cydia pomonella* L.) (Lepidoptera: Tortricidae). *Biocontrol* 55, 777-788.
12. **Salman, M. (2010):** Determination of antibiotic activity on plasmids from fluorescent pseudomonads isolates CW2, WB15 and WB52 against pre-emergence damping-off caused by *Pythium ultimum* and *Rhizoctonia solani* in cucumber. *Biological Control* 53, 161-167.
13. **Sun, C.A., Johnson, J.M., Cai, D.G., Sherameti, I., Oelmüller, R. & Lou, B.G. (2010):** *Piriformospora indica* confers drought tolerance in Chinese cabbage leaves by stimulating antioxidant enzymes, the expression of drought-related genes and the plastid-localized CAS protein. *Plant Physiology* 167, 1009-1017.
14. **Thurau, T., Ye, W.Z., Menkhaus, J., Knecht, K., Tang, G. & Cai, D.G. (2010):** Plant nematode control. *Sugar Technology* 12, 229-237.
15. **Toepfer, S., Burger, R., Ehlers, R.-U., Peters, A. & Kuhlmann, U. (2010):** Controlling western corn rootworm larvae with entomopathogenic nematodes: effect of application techniques on plant-scale efficacy. *Journal of Applied Entomology* 134, 467-480.

16. **Toepfer, S., Hatala-Zseller, I., Ehlers, R.-U., Peters, A. & Kuhlmann, U. (2010):** The effect of application techniques on field-scale efficacy: can the use of entomopathogenic nematodes reduce damage by western corn rootworm larvae? *Agricultural and Forest Entomology* 12, 389-402.
17. **Wang, Y., Shen, D., Bo, S.P., Chen, H.A., Zheng, J.A., Zhu, Q.-H., Cai, D.G., Helliwell, C. & Fan, L.J. (2010):** Sequence variation and selection of small RNAs in domesticated rice. *BMC Evolutionary Biology* 10, 119-125.

2009

Bücher/Buchkapitel:

1. **Börner, H., Schlüter, K. & Aumann, J. (2009):** Pflanzenkrankheiten und Pflanzenschutz, 8. Auflage. Springer, Dordrecht, Heidelberg, London, New York.

Begutachtete Publikationen:

1. **Abd-El Salam, K.A., Guo, J.R., Moslem, M.A., Bahkali, A.H. & Verreet, J.-A. (2009):** Suitability of intergenic spacer or internal transcribed spacer microsatellite-primed PCR for the identification of *Rhizoctonia solani* and some phytofungi. *Journal of Rapid Methods and Automation in Microbiology* 17, 383-397.
2. **Endrigkeit, J., Wang, X.X., Cai, D.G., Zhang, C.Y., Long, Y., Meng, J.L. & Jung, C. (2009):** Genetic mapping, cloning and functional characterization of the BnaX.VTE4 gene encoding a γ -tocopherol methyltransferase from oilseed rape. *Theoretical and Applied Genetics* 119, 567-575.
3. **Hirao, A. & Ehlers, R.-U. (2009):** Effect of temperature on the development of *Steinernema carpocapsae* and *Steinernema feltiae* (Nematoda: Rhabditida) in liquid culture. *Applied Microbiology and Biotechnology* 84, 1061-1067.
4. **Hirao, A. & Ehlers, R.-U. (2009):** Influence of cell density and phase variants of bacterial symbionts (*Xenorhabdus* spp.) on dauer juvenile recovery and development of biocontrol nematodes *Steinernema carpocapsae* and *S. feltiae* (Nematoda: Rhabditida). *Applied Microbiology and Biotechnology* 84, 77-85.
5. **Knott, J.M. (2009):** Biosynthesis of long-chain polyamines by crenarchaeal polyamine synthases from *Hyperthermus butylicus* and *Pyrobaculum aerophilum*. *FEBS Letters* 583, 3519-3524.
6. **Kurtz, B., Hitpold, I., Turlings, T.C.J., Kuhlmann, U. & Toepfer, S. (2009):** Comparative susceptibility of larval instars and pupae of the western corn rootworm to infection by three entomopathogenic nematodes. *Biocontrol* 54, 255-262.
7. **Scheider, N., Guo, J.R., Verreet, J.-A. & Beyer, M. (2009):** Assessing the intensity of *Fusarium*-damage in wheat: A comparison of selected disease parameters during disease development and the role of fungicides. *Journal of Plant Diseases and Protection* 116, 118-123.
8. **Schlagenhafer, S., Engelhard, B. & Wolf, P.F.J. (2009):** Sporenverteilung des Echten Mehltaus im Hopfen ausgehend von Infektionsherden und Abhängigkeit vom Vegetationsstadium. *Gesunde Pflanzen* 61, 31-37.
9. **Toepfer, S., Haye, T., Erlandson, M., Goettel, M., Lundgren, J.G., Kleespies, R.G., Weber, D.C., Walsh, G.C., Peters, A., Ehlers, R.-U., Strasser, H., Moore, D., Keller, S., Vidal, S. & Kuhlmann, U. (2009):** A review of the natural enemies of beetles in the subtribe Diabroticina (Coleoptera: Chrysomelidae): implications for sustainable pest management. *Biocontrol Science and Technology* 19, 1-65.
10. **Wang, Y., Yu, T., Li, Y., Cai, D.G., Liu, X., Lu, H. & Zheng, X.D. (2009):** Postharvest biocontrol of *Alternaria alternata* in Chinese winter jujube by *Rhodosporidium paludigenum*. *Journal of Applied Microbiology* 107, 1492-1498.
11. **Wolf, P.F.J., Lenz, R., Baron, K. & Verreet, J.-A. (2009):** Quaternäres Integriertes Pflanzenschutz-System für den Echten Mehltau der Zuckerrübe. III.

Ökonomische Schadensschwelle und Verlustprognose. Journal für Kulturpflanzen 61, 91-97.

12. **Wolf, P.F.J. & Verreet, J.-A. (2009):** Empirisch-deterministische Prognose von Befall und Verlust verursacht durch die *Cercospora*-Blattfleckenkrankheit in Zuckerrüben. Journal für Kulturpflanzen 61, 168-177.

2008

Bücher/Buchkapitel:

1. **Hauschildt, R., Speiser, B., Tamm, L. & Ehlers, R.-U. (2008):** Réglementation et homologation des produits phyto-pharmaceutiques à base d`extraits végétaux dans la législation de l` Union européenne (UE) : présent et perspectives d` évolution. In : C. Regnault-Roger, B.J.R. Philogène, C. Vincent (eds.): Biopesticides d` origine végétale, 477-496. Lavoisier, Paris.

Begutachtete Publikationen:

1. **Abd-Elsalam, K.A., Bahkali, H.A., Al-Khedhairi, A.A. & Verreet, J.-A. (2008):** Development of a conventional and Lightcycler PCR assay for detection of *Fusarium solani*. Gene Genomes Genomics 2, 63–67.
2. **Beyer, M. & Aumann, J. (2008):** Effects of *Fusarium* infection on the amino acid composition of winter wheat grain. Food Chemistry 111, 759-754.
3. **Binggan, L., Chao S. & Cai, D.G. (2008):** *Piriformospora indica*, a growth promoting fungus with multiple biotechnological applications. Acta Phytophylactica Sinica 34, 653-656.
4. **Guo, J.R. & Verreet, J.-A. (2008):** Formation and germination of *Septoria tritici* secondary conidia as affected by environmental factors. Journal of Phytopathology 156, 635-637.
5. **Guo, X., Ruan, S., Hu, W., Cai, D.G. & Fan, L. (2008)** Chloroplast DNA insertions into the nuclear genome of rice: the genes, sites and ages of insertion involved. Functional and Integrative Genomics 8, 101-108.
6. **Hirao, A. & Ehlers, R.-U. (2008):** Influence of nematode inoculum density and temperature on development of *Steinernema carpocapsae* and *S. feltiae* in liquid culture. Communications in Agricultural and Applied Biological Sciences 73 (4), 699-702.
7. **Klix, M.B., Beyer, M. & Verreet, J.-A. (2008):** Effects of cultivar, agronomic practices, geographic location, and meteorological conditions on the composition of selected *Fusarium* species on wheat heads. Canadian Journal of Plant Pathology 30, 46-57.
8. **Mukuka, J., Strauch, O. & Ehlers, R.-U. (2008):** Variability in desiccation toerance among different strains oft he entomopathogenic nematode *Heterorhabditis bacteriophora*. Communications in Agricultural and Applied Biological Sciences 73 (4), 669-672.
9. **Susurluk, A. & Ehlers, R.-U. (2008):** Field persistence of the entomopathogenic nematode *Heterorhabditis bacteriophora* in different crops. Biocontrol 53, 627-641.
10. **Susurluk, A. & Ehlers, R.-U. (2008):** Sustainable control of black vine weevil larvae, *Otiorhynchus sulcatus* (Coleoptera: Curculionidae) with *Heterorhabditis bacteriophora* in strawberry. Biocontrol Science and Technology 18, 635-640.
11. **Toepfer, S., Peters, A., Ehlers, R.-U. & Kuhlmann, U. (2008):** Comparative assessment of the efficacy of entomopathogenic nematode species at reducing western corn rootworm larvae and root damage in maize. Journal of Applied Entomology 132, 337-348.
12. **Wolf, P.F.J. & Verreet, J.-A. (2008):** Quaternary IPM (integrated pest management) – concept for the control of powdery mildew in sugar beets. Communications in Agricultural and Applied Biological Sciences 73 (2), 57-68.

2007

Bücher/Buchkapitel:

1. **Ehlers, R.-U. (2007):** Entomopathogenic nematodes: From science to commercial use. In: C. Vincent, M.S. Goettel, G. Lazarovits (eds.): Biological Control - A Global Perspective: Case Studies from Around the World, 136-151. CABI Publishing, Wallingford.

Begutachtete Publikationen:

1. **Beyer, M., Klix, M.B. & Verreet, J.-A. (2007):** Estimating mycotoxin contents of *Fusarium*-damaged winter wheat kernels. International Journal of Food Microbiology 119, 153-158.
2. **Bremer, H. (2007):** Entwicklung einer Herbstprognose für den Erreger der Wurzelhals- und Stängelfaule *Phoma lingam* (Teleomorph: *Leptosphaeria maculans*). Gesunde Pflanzen 59, 161-169.
3. **Dantigny, P., Marin, S., Beyer, M. & Magan, N. (2007):** Mould germination: Data treatment and modelling. International Journal of Food Microbiology 114, 17-24.
4. **Guo, J.R., Schnieder, F. & Verreet, J.-A. (2007):** A real-time PCR assay for quantitative and accurate assessment of fungicide effects on *Mycosphaerella graminicola* leaf blotch. Journal of Phytopathology 155, 482-487.
5. **Henze, M., Beyer, M., Klink, H. & Verreet, J.-A. (2007):** Characterizing meteorological scenarios favorable for *Septoria tritici* infections in wheat and estimation of latent periods. Plant Disease 91, 1445-1449.
6. **Klix, M.B., Verreet, J.-A. & Beyer, M. (2007):** Comparison of the declining triazole sensitivity of *Gibberella zeae* and increased sensitivity achieved by advances in triazole fungicide development. Crop Protection 26, 683-690.
7. **Krome, K., Kabsch, U. & Aumann, J. (2007):** The effect of benzothiadiazole and fungal extracts of *Cercospora beticola* and *Fusarium graminearum* on phosphoenolpyruvate carboxylase activity in cucumber leaves. Journal of Plant Diseases and Protection 114, 250-255.
8. **Kurtz, B., Toepfer, S., Ehlers, R.-U. & Kuhlmann, U. (2007):** Assessment of establishment and persistence of entomopathogenic nematodes for biological control of western corn rootworm. Journal of Applied Entomology 131, 420-425.
9. **Lein, J.C., Asbach, K., Tian, Y.Y., Schulte, D., Li, C.Y., Koch, G., Jung, C. & Cai, D.G. (2007):** Resistance gene analogues are clustered on chromosome 3 of sugar beet and cosegregate with QTL for rhizomania resistance. Genome 50, 61-71.
10. **Lou, B.G., Sun, C.A. & Cai, D.G. (2007):** *Piriformospora indica*, a growth promoting fungus with multiple biotechnological applications. Acta Phytophylactica Sinica 34, 653-656.
11. **Oestergaard, J., Ehlers, R.-U., Martinez-Ramirez, A.C. & Real, M.D. (2007):** Binding of Cyt1Aa and Cry11Aa toxins of *Bacillus thuringiensis* serovar *israelensis* to brush border membrane vesicles of *Tipula paludosa* (Diptera: Nematocera) and subsequent pore formation. Applied and Environmental Microbiology 73, 3623-3629.
12. **Oestergaard, J., Voss, S., Lange, H., Lemke, H., Strauch, O. & Ehlers, R.-U. (2007):** Quality control of *Bacillus thuringiensis* ssp. *israelensis* products based on toxin quantification with monoclonal antibodies. Biocontrol Science and Technology 17, 295-302.
13. **Omar, M.R., Abd-Elsalam, K.A., Aly, A.A., El-Samawaty, A.M.A. & Verreet, J.-A. (2007):** Diversity of *Macrophomina phaseolina* from cotton in Egypt: Analysis of pathogenicity, chlorate phenotypes, and molecular characterization. Journal of Plant Diseases and Protection 114, 196-204.

14. **Ragab, W.S.M., Drusch, S., Schnieder, F. & Beyer, M. (2007):** Fate of deoxynivalenol in contaminated wheat grain during preparation of Egyptian 'balila'. *International Journal of Food Sciences and Nutrition* 58, 169-177.
15. **Unlo, I.O., Ehlers, R.-U. & Susurluk, A. (2007):** Additional data and first record of the entomopathogenic nematode *Steinernema weiseri* from Turkey. *Nematology* 9, 739-741.
16. **Wolf, P.F.J., Lenz, R., Baron, K. & Verreet, J.-A. (2007):** Quaternäres Integriertes Pflanzenschutz-System für den Echten Mehltau der Zuckerrübe. II. Prüfung von epidemieorientierten Bekämpfungsschwellen als Indikatoren des Fungizideinsatzes. *Nachrichtenblatt des Deutschen Pflanzenschutzdienstes* 59, 107-112.

2006

Bücher/Buchkapitel:

1. **Ehlers, R.-U. (2006):** Potenziale des biologischen Pflanzenschutzes. In: A. von Tiedemann, F. Feldmann (eds.): *Biotechnologie - Innovationsmotor für den Pflanzenschutz*, 17-32. Eugen Ulmer, Stuttgart.
2. **Melo, E. L., C. A. Ortega, Gaigl, A., Ehlers, R.-U. & Bellotti, A. (2006):** Parasitismo de dos cepas de entomonematodos, una nativa (*Steinernema feltiae*, cepa Colombia) y otra introducción (*Heterorhabditis bacteriophora* cepa E-Nema), sobre los seis estados de desarrollo de *Cyrtomenus bergi* Froeschner (Heteroptera: Cydnidae) en condiciones de laboratorio. In: J.C. Parada S., J.E. Luque Z., W. de J. Piedrahíta C. (eds.): *Nematodos Entomoparásitos: Experiencias y Perspectivas*, 175-183. Universidad Nacional de Colombia, Bogotá.
3. **Melo, E.L., Ortega, C.A., Susurluk, A., Gaigl, A. & Ehlers, R.-U. (2006):** Búsqueda de nematodos entomoparásitos asociados a insectos rizófagos en regiones de Colombia y Panamá. In: . C. Parada S., J.E. Luque Z., W. de J. Piedrahíta C. (eds.): *Nematodos Entomoparásitos: Experiencias y Perspectivas*, 28-32. Universidad Nacional de Colombia, Bogotá.

Begutachtete Publikationen:

1. **Beyer, M., Klix, M.B., Klink, H. & Verreet, J.-A. (2006):** Quantifying the effects of previous crop, tillage, cultivar and triazole fungicides on the deoxynivalenol content of wheat grain – a review. *Journal of Plant Diseases and Protection* 113, 241-246.
2. **Ciche, T.A., Darby, C., Ehlers, R.-U., Forst, S. & Goodrich-Blair, H. (2006):** Dangerous liaisons: The symbiosis of entomopathogenic nematodes and bacteria. *Biological Control* 38, 22-46.
3. **Elsa, L.M.M., Ortega-Ojeda, C.A., Gaigl, A., Ehlers, R.-U. & Bellotti, A.C. (2006):** Evaluation of two commercial strains of entomonematodes as control agents of *Cyrtomenus bergi* Froeschner (Hemiptera: Cydnidae). *Revista Colombiana de Entomología* 32, 31-38.
4. **Guo, J.R., Schnieder, F. & Verreet, J.-A. (2006):** Presymptomatic and quantitative detection of *Mycosphaerella graminicola* development in wheat using a real-time PCR assay. *FEMS Microbiology Letters* 262, 223-229.
5. **Guo, J.R., Schnieder, F. & Verreet, J.-A. (2006):** Differences between the fingerprints generated from total RNA and poly-A RNA using a modified procedure of cDNA-AFLP and silver staining. *Biotechnology Letters* 28, 267-270.
6. **Kaya, H.K., Aguilera, M.M., Alumai, A., Choo, H.Y., de la Torre, M., Fodor, A., Ganguly, S., Hazir, S., Lakatos, T., Pye, A., Wilson, M., Yamanaka, S., Yang, H.W. & Ehlers, R.-U. (2006):** Status of entomopathogenic nematodes and their symbiotic bacteria from selected countries or regions of the world. *Biological Control* 38, 134-155.

7. **Klix, M.B., Beyer, M. & Verreet, J.-A. (2006):** Declining triazole sensitivity in *Gibberella zeae* and advances in fungicide development. *Communications in Agricultural and Applied Biological Sciences* 71 (3a), 849-852.
8. **Oestergaard, J., Belau, C., Strauch, O., Ester, A., van Rozen, K. & Ehlers, R.-U. (2006):** Biological control of *Tipula paludosa* (Diptera: Nematocera) using entomopathogenic nematodes (*Steinernema* spp.) and *Bacillus thuringiensis* subsp. *israelensis*. *Biological Control* 39, 525-531.
9. **Susurluk, I.A. & Toprak, U. (2006):** Molecular identification of three entomopathogenic nematodes from Turkey by PCR-RFLP of the ITS regions. *Phytoparasitica* 34, 17-20.
10. **Verreet, J.-A. & Aumann, J. (2006):** Preface by the editor. *Journal of Plant Diseases and Protection* 113, 1.
11. **Wieczorek, K., Golecki, B., Gerdes, L., Heinen, P., Szakasits, D., Durachko, D.M., Cosgrove, D.J., Kreil, D.P., Puzio, P.S., Bohlmann, H. & Grundler, F.M.W. (2006):** Expansis are involved in the formation of nematode-induced syncytia in roots of *Arabidopsis thaliana*. *Plant Journal* 48, 98-112.
12. **Wolf, P.F.J., Lenz, R., Baron, K. & Verreet, J.-A. (2006):** Quaternary integrated pest management concept for powdery mildew in sugar beet. I. Analysis of epidemic determinants to predict disease onset. *Journal of Plant Diseases and Protection* 113, 61-67.
13. **Yi, X. & Ehlers, R.-U. (2006):** Combining *Steinernema carpocapsae* and *Bacillus thuringiensis* strains for control of diamondback moth (*Plutella xylostella*). *Communications in Agricultural and Applied Biological Sciences* 71 (3a), 633-636.

2005

1. **Beyer, M., Verreet, J.-A. & Ragab, W.S.M. (2005):** Effect of relative humidity on germination of ascospores and macroconidia of *Gibberella zeae* and deoxynivalenol production. *International Journal of Food Microbiology* 98, 233-240.
2. **Beyer, M. & Verreet, J.-A. (2005):** Germination of *Gibberella zeae* ascospores as affected by age of spores after discharge and environmental factors. *European Journal of Plant Pathology* 111, 381-389.
3. **Cowles, R.S., S. Polavarapu, Williams, R.N., Thies, A. & Ehlers, R.-U. (2005):** Soft fruit applications. In: P.S. Grewal, R.-U. Ehlers, D.I. Shapiro-Ilan (eds.): *Nematodes as Biocontrol Agents*, 231-254. CABI Publishing, Wallingford.
4. **Drusch, S. & Aumann, J. (2005):** Mycotoxins in fruits: Microbiology, occurrence, and changes during fruit processing. In: S.L. Taylor (ed.): *Advances in Food and Nutrition Research*, Vol. 50, 33-78. Academic Press, San Diego, London.
5. **Ehlers, R.-U. (2005):** Forum on safety and regulation. In: P.S. Grewal, R.-U. Ehlers, D.I. Shapiro-Ilan (eds.): *Nematodes as Biocontrol Agents*, 107-114. CABI Publishing, Wallingford.
6. **Ehlers, R.-U., Oestergaard, J., Hollmer, S., Wingen, M. & Strauch, O. (2005):** Genetic selection for heat tolerance and low temperature activity of the entomopathogenic nematode-bacterium complex *Heterorhabditis bacteriophora* – *Photorhabdus luminescens*. *Biocontrol* 50, 699-716.
7. **Ehlers, R.-U. & Shapiro-Ilan, D. I. (2005):** Mass production. In: P.S. Grewal, R.-U. Ehlers, D.I. Shapiro-Ilan (eds.): *Nematodes as Biocontrol Agents*, 65-78. CABI Publishing, Wallingford.
8. **Grewal, P.S., Ehlers, R.-U. & Shapiro-Ilan, D.I., eds. (2005):** *Nematodes as Biocontrol Agents*. CABI Publishing, Wallingford.
9. **Grewal, P.S., Ehlers, R.-U. & Shapiro-Ilan, D. (2005):** Critical issues and research needs for expanding the use of nematodes in Biocontrol. In: P.S. Grewal,

- R.-U. Ehlers, D.I. Shapiro-Ilan (eds.): Nematodes as Biological Agents, 479-489. CABI Publishing, Wallingford.
10. **Guo, J.R., Schnieder, F., Abd-Elsalam, K.A. & Verreet, J.-A. (2005):** Rapid and efficient extraction of genomic DNA from different phytopathogenic fungi using DNAzol reagent. *Biotechnology Letters* 27, 3-6.
 11. **Guo, J.R., Schnieder, F., Beyer, M. & Verreet, J.-A. (2005):** Rapid detection of *Mycosphaerella graminicola* in wheat using reverse transcription-PCR assay. *Journal of Phytopathology* 153, 674-679.
 12. **Kruse, T. & Verreet, J.-A. (2005):** Epidemiological studies on winter oilseed rape (*Brassica napus* L. var. *napus*) infected by *Phoma lingam* (teleomorph *Leptosphaeria maculans*) and the effects of different fungicide applications with Folicur® (tebuconazole). *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz* 112, 17-41.
 13. **Ludewig, A., Kabsch, U. & Verreet, J.-A. (2005):** Mycotoxin production of 31 *Fusarium graminearum* isolates and aggressiveness against wheat seedlings. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz* 112, 124-133.
 14. **Ludewig, A., Kabsch, U. & Verreet, J.-A. (2005):** Comparative deoxynivalenol accumulation and aggressiveness of isolates of *Fusarium graminearum* on wheat and the influence on yield as affected by fungal isolate and wheat cultivar. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz* 112, 329-342.
 15. **Ragab, W.S.M., Drusch, S., Kuhlmann, A. & Beyer, M. (2005):** Characterizing effects of fermentation and baking on the deoxynivalenol content of rolls. *Journal of Applied Botany and Food Quality* 79, 197-201.
 16. **Schroer, S. & Ehlers, R.-U. (2005):** Foliar application of the entomopathogenic nematode *Steinernema carpocapsae* for biological control of diamondback moth larvae (*Plutella xylostella*). *Biological Control* 33, 81-86.
 17. **Schroer, S., Sulistyanto, D. & Ehlers, R.-U. (2005):** Control of *Plutella xylostella* using polymer-formulated *Steinernema carpocapsae* and *Bacillus thuringiensis* in cabbage fields. *Journal of Applied Entomology* 129, 198-204.
 18. **Schroer, S., Xi, X.L. & Ehlers, R.-U. (2005):** Evaluation of adjuvants for foliar application of *Steinernema carpocapsae* against larvae of the diamondback moth (*Plutella xylostella*). *Nematology* 7, 37-44.
 19. **Schroer, S., Ziermann, D. & Ehlers, R.-U. (2005):** Mode of action of a surfactant-polymer formulation to support performance of the entomopathogenic nematode *Steinernema carpocapsae* for control of diamondback moth larvae (*Plutella xylostella*). *Biocontrol Science and Technology* 15, 601-613.
 20. **Toepfer, S., Gueldenzoph, C., Ehlers, R.-U. & Kuhlmann, U. (2005):** Screening of entomopathogenic nematodes for virulence against the invasive western corn rootworm, *Diabrotica virgifera virgifera* (Coleoptera: Chrysomelidae) in Europe. *Bulletin of Entomological Research* 95, 473-482.
 21. **Wolf, P.F.J. & Verreet, J.-A. (2005):** Factors affecting the onset of *Cercospora* leaf spot epidemics in sugar beet and establishment of disease-monitoring thresholds. *Phytopathology* 95, 269-274.

2004

1. **Beyer, M., Röding, S., Ludewig, A. & Verreet, J.-A. (2004):** Germination and survival of *Fusarium graminearum* macroconidia as affected by environmental factors. *Journal of Phytopathology* 152, 92-97.
2. **Hazir, S., Stackebrandt, E., Lang, E., Schumann, P., Ehlers, R.-U. & Keskin, N. (2004):** Two new subspecies of *Photorhabdus luminescens*, isolated from *Heterorhabditis bacteriophora* (Nematoda: Heterorhabditidae): *Photorhabdus luminescens* subsp. *kayaii* subsp. nov. and *Photorhabdus luminescens* subsp. *thracensis* subsp. nov. *Systematic and Applied Microbiology* 27, 36-42.

3. **Johnigk, S.A., Ecke, F., Poehling, H.-M. & Ehlers, R.-U. (2004):** Liquid culture mass production of biocontrol nematodes, *Heterorhabditis bacteriophora* (Nematoda: Rhabditida): improved timing of dauer juvenile inoculation. *Applied Microbiology and Biotechnology* 64, 651-658.
4. **Karanastasi, E., Decraemer, W., Wyss, U. & Brown, D.J.K. (2004):** The ultrastructure of the feeding apparatus and pharyngeal tract of four European species of Trichodoridae (Nematoda: Triplonchida). *Nematology* 6, 695-713.
5. **Strauch, O., Oestergaard, J., Hollmer, S. & Ehlers, R.-U. (2004):** Genetic improvement of the desiccation tolerance of the entomopathogenic nematode *Heterorhabditis bacteriophora* through selective breeding. *Biological Control* 32, 218-226.
6. **Struck, E., Ebssa, L., Ehlers, R.-U., Poehling, H.-M., Gaigl, A. & Borgemeister, C. (2004):** Interactions between host plants, the subterranean burrower bug, *Cyrtomenus bergi*, and the entomopathogenic nematode *Heterorhabditis megidis*. *Nematology* 6, 633-639.
7. **Wolf, P.F.J., Klink, H. & Verreet, J.-A. (2004):** A concept in integrated pest management (IPM) of fungal leaf diseases of cereals and sugar beets. In: R.T. Lartey, J.A. Caesar (eds.): *Emerging Concepts in Plant Health Management*, 69-102. Research Signpost, Trivandrum, Kerala.

2003

1. **Abd-Elsalam, K.A., Schnieder, F., Asran-Amal, M.A., Khalil, M.S. & Verreet, J.-A. (2003):** Intraspecies genomic groups in *Fusarium semitectum* and their correlation with origin and morphological characteristics. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz* 110, 409-418.
2. **Abd-Elsalam, K.A., Schnieder, F., Khalil, M.S., Aly, A.A. & Verreet, J.-A. (2003):** Genetic variation at the intra- and interspecific level in *Fusarium* spp. associated from Egyptian cottons. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz* 110, 46-53.
3. **Abd-Elsalam, K.A., Schnieder, F., Khalil, M.S., Asran-Amal, M.A. & Verreet, J.-A. (2003):** Use of AFLP fingerprinting to analyze genetic variation within and between populations of *Fusarium* spp. derived from Egyptian cottons. *Journal of Plant Pathology* 85, 99-103.
4. **Ehlers, R.-U. (2003):** Entomopathogenic nematodes in the European biocontrol market. *Communications in Agricultural and Applied Biological Sciences* 68 (4a), 3-16.
5. **Ehlers, R.-U. (2003):** Biocontrol nematodes. In: H.M.T. Hokkanen, A.E. Hajek (eds.): *Environmental Impacts of Microbial Insecticides*, 177-220. Kluwer, Dordrecht.
6. **Juergensen, K., Scholz-Starke, J., Sauer, N., Hess, P., van Bel, A.J.E. & Grundler, F.M.W. (2003):** The companion cell-specific Arabidopsis disaccharide carrier AtSUC2 is expressed in nematode-induced syncytia. *Plant Physiology* 131, 61-69.
7. **Karanastasi, E., Wyss, U. & Brown, D.J.K. (2003):** An *in vitro* examination of the feeding behaviour of *Paratrichodorus anemones* (Nematoda: Trichodoridae), with comments on the ability of the nematode to acquire and transmit tobnavirus particles. *Nematology* 5, 421-434.
8. **Mölick, G. & Wyss, U. (2003):** Effect of aphid-infested plants on the host location and learning behaviour of the parasitoid *Aphelinus abdominalis*. *Communications in Agricultural and Applied Biological Sciences* 68(4b), 167-177.
9. **Premachandra, W.T.S.D., Borgemeister, C., Berndt, O., Ehlers, R.-U. & Poehling, H.-M. (2003):** Combined releases of entomopathogenic nematodes and the predatory mite *Hypoaspis aculeifer* to control soil-dwelling stages of western flower thrips *Frankliniella occidentalis*. *Biocontrol* 48, 529-541.

10. **Premachandra, W.T.S.D., Borgemeister, C., Berndt, O., Ehlers, R.-U. & Poehling, H.-M. (2003):** Laboratory bioassays of virulence of entomopathogenic nematodes against soil-inhabiting stages of *Frankliniella occidentalis* Pergande (Thysanoptera: Thripidae). *Nematology* 5, 539-547.
11. **Rühm, R., Dietsche, E., Harloff, H.-J., Lieb, M., Franke, S. & Aumann, J. (2003):** Characterization and partial purification of a white mustard kairomone that attracts the beet cyst nematode, *Heterodera schachtii*. *Nematology* 5, 17-22.
12. **Verreet, J.-A., Heger, M., Oerke, E.-C., Dehne, H.-W., Finger, I., Busse, C. & Klink, H. (2003):** The IPM wheat model – results of a three-year study in North Rhine-Westphalia, Lower Saxony and Schleswig-Holstein. *Communications in Agricultural and Applied Biological Sciences* 68 (4b), 499-509.
13. **Wolf, P.F.J. & Verreet, J.-A. (2003):** Innovative combination of IPM (integrated pest management) tools – The IPM sugar beet model. *Communications in Agricultural and Applied Biological Sciences* 68 (4b), 491-498.

2002

1. **Bayer, K., Mölck, G., Petersen, G. & Wyss, U. (2002):** Eigenschaften des weiblichen Sexualpheromons des Blattlausparasitoiden *Aphidius uzbekistanicus* und seine Bedeutung für die Partnerfindung. *Mitteilungen der Deutschen Gesellschaft für Allgemeine und Angewandte Entomologie* 13, 393- 397.
2. **Bochow, H., Koch, K.-H., Poehling, H.-M., von Alten, H. & Wyss, U. (2002):** Mechanisms of interactions in the system plant, parasite and beneficial organisms. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz* 108, 626- 652.
3. **Borgemeister, C., Ebssa, L., Premachandra, D., Berndt, O., Ehlers, R.-U. & Poehling, H.-M. (2002):** Biological control of soil-dwelling life stages of western flower thrips *Frankliniella occidentalis* (Pergande) (Thysanoptera: Thripidae) by entomopathogenic nematodes and *Hypoaspis* spp. (Acari: Laelapidae). *IOBC/WPRS Bulletin* 25, 29-32.
4. **Charwat, S.M., Fisher, J.M. & Wyss, U. (2002):** The effect of osmotic stress on desiccation survival and water content of four nematode species. *Nematology* 4, 89-97.
5. **Hass, B., Downes, M.J. & Griffin, C.T. (2002):** Persistence of four *Heterorhabditis* spp. isolates in soil: Role of lipid reserves. *Journal of Nematology* 34, 151-158.
6. **Johnigk, S.-A., Hollmer, S., Strauch, O., Wyss, U. & Ehlers, R.-U. (2002):** Heritability of the liquid culture mass production potential of the entomopathogenic nematode *Heterorhabditis bacteriophora*. *Biocontrol Science and Technology* 12, 267-276.
7. **Mölck, G. & Wyss, U. (2002):** *Aphelinus abdominalis*, ein erfolgversprechender Nützling zur Blattlausbekämpfung. *Gesunde Pflanzen* 54, 92-97.
8. **Petersen, G., Matthiesen, C., Stolzenburg, N., Zimmermann, N., Hillgraf, R., Lehmann, L., Franke, W. & Wyss, U. (2002):** Die verhaltensmodifizierende Wirkung von *Alloxysta-victrix*-Duftstoffen im System Blattlaus-Primärparasitoid-Hyperparasitoid. *Mitteilungen der Deutschen Gesellschaft für Allgemeine und Angewandte Entomologie* 13, 51-55.
9. **Wyss, U. (2002):** Feeding behaviour of plant-parasitic nematodes. In: D. Lee (ed.): *Biology of Nematodes*, 233-259. Taylor and Francis, London.

2001

1. **Aumann, J. & Ehlers, R.-U. (2001):** Physico-chemical properties and mode of action of a signal from the symbiotic bacterium *Photorhabdus luminescens* inducing

dauer juvenile recovery in the entomopathogenic nematode *Heterorhabditis bacteriophora*. *Nematology* 3, 849-853.

2. **Boemare, N., Ehlers, R.-U., Burnell, A.M., Wegensteiner, R., Koschier, E. & Mulder, R. (2001):** COST Action 819 Entomopathogenic nematodes - Virulence factors and secondary metabolites from symbiotic bacteria of entomopathogenic nematodes. Office for Official Publications of the EC, Luxembourg.
3. **Cai, D.G., Wyss, U., Jung, C. & Kleine, M. (2001):** Genetic Engineering for resistance to nematodes. In: G. Khachatourians, A. McHughen, R. Scorza, W.K. Nip, J.K. Hui (eds.): *Transgenic Plants and Crops*, 233-248. Marcel Dekker, New York.
4. **Ehlers, R.-U. (2001):** Mass production of entomopathogenic nematodes for plant protection. *Applied Microbiology and Biotechnology* 56, 623-633.
5. **Han, R. & Ehlers, R.-U. (2001):** Effect of *Photorhabdus luminescens* phase variants on the *in vivo* and *in vitro* development and reproduction of *Heterorhabditis bacteriophora* and *Steinernema carpocapsae*. *FEMS Microbiology Ecology* 35, 239-247.
6. **Han, R. & Ehlers, R.-U. (2001):** *In vivo* and *in vitro* reproduction of *Steinernema carpocapsae* and *Heterorhabditis bacteriophora* on *Photorhabdus luminescens* phase variants. In: N. Boemare, R.-U. Ehlers, A. M. Burnell, R. Wegensteiner, E. Koschier, R. Mulder (eds.): *COST 819 Entomopathogenic nematodes - Virulence factors and secondary metabolites from symbiotic bacteria of entomopathogenic nematodes*, 33-45. Office for Official Publications of the EC, Luxembourg.
7. **Han, R. & Ehlers, R.-U. (2001):** Nematicidal activity of culture filtrates from *Photorhabdus luminescens*. In: N. Boemare, R.-U. Ehlers, A. M. Burnell, R. Wegensteiner, E. Koschier, R. Mulder (eds.): *COST 819 Entomopathogenic nematodes - Virulence factors and secondary metabolites from symbiotic bacteria of entomopathogenic nematodes*, 109-119. Office for Official Publications of the EC, Luxembourg.
8. **Hussein, M. & Ehlers, R.-U. (2001):** Significance of the *Photorhabdus luminescens* inclusion protein for the development of *Heterorhabditis bacteriophora*. In: N. Boemare, R.-U. Ehlers, A. M. Burnell, R. Wegensteiner, E. Koschier, R. Mulder (eds.) *COST 819 Entomopathogenic nematodes - Virulence factors and secondary metabolites from symbiotic bacteria of entomopathogenic nematodes*, 47-53. Office for Official Publications of the EC, Luxembourg.
9. **Mölk, G. & Wyss, U. (2001):** The effect of experience on the parasitization efficiency of the aphid antagonist *Aphelinus abdominalis* in greenhouse crops. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz* 108, 616-625.
10. **Nitz, I., Berkefeld, H., Puzio, P.S. & Grundler, F.M.W. (2001):** *Pyk 10*, a seedling and root specific gene and promoter from *Arabidopsis thaliana*. *Plant Science* 161, 337- 346.
11. **Poehling, H.-M., von Alten, H. & Wyss, U. (2001):** Mechanisms of interaction in the system plant, parasite and beneficial organisms. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz* 108, 626-652.
12. **Susurluk, A., Dix, I., Stackebrandt, E., Strauch, O., Wyss, U. & Ehlers, R.-U. (2001):** Identification and ecological characterisation of three entomopathogenic nematode-bacterium complexes from Turkey. *Nematology* 3, 833-841.
13. **Vierheilig, H., Knoblauch, M., Juergensen, K., van Bel, A., Grundler, F.M.W. & Piché, Y. (2001):** Imaging arbuscular mycorrhizal structures in living roots of *Nicotiana tabacum* by light, epifluorescence, and confocal laser scanning microscopy. *Canadian Journal of Botany* 79, 231-237.
14. **Vierheilig, H., Maier, W., Wyss, U., Samson, J., Strack, D. & Piché, Y. (2001):** Cyclohexenone derivative- and phosphate-levels in split-root systems and their role in the systemic suppression of mycorrhization in precolonized barley plants. *Journal of Plant Physiology* 157, 593-599.

15. **Wolf, P.F.J., Heindl, M. & Verreet, J.-A. (2001):** Zum Einfluß des Bestandesklimas auf die Prädisposition der Zuckerrübe gegenüber Infektionen von *Cercospora beticola* (Sacc.). Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz 108, 578-592.
16. **Wolf, P.F.J., Weis, F.J. & Verreet, J.-A. (2001):** Bekämpfungsschwellen als Indikatoren des Fungizideinsatzes zur Kontrolle des Blattbefalls von *Cercospora beticola* (Sacc.) in Zuckerrüben. Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz 108, 244-257.
17. **Wyss, U. & Poehling, H.-M. (2001):** Multitrophic interactions between plants, herbivores and their carnivorous natural enemies mediated by infochemicals. Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz 108, 630 – 642.

2000

1. **Ehlers, R.-U., Niemann, I., Hollmer, S., Strauch, O., Jende, D., Shanmugasundaram, M., Mehta, U.K., Easwaramoorthy, S.K. & Burnell, A. (2000):** Mass production potential of the bacto-helminthic biocontrol complex *Heterorhabditis indica* - *Photorhabdus luminescens*. Biocontrol Science and Technology 10, 607-616.
2. **Gheysen, G., Abad, P., Bleve, T., Blok, V., Fenoll, C., Gatehouse, J., Grundler, F.M.W., Lindsey, K., Ohl, S., Sagen, K., Shields, R. & Helder, H. (2000):** Concerted efforts to develop handles for plant parasitic nematode control. In: G.E. de Vries, K. Metzloff (eds.): Phytosfere'99 - Highlights in European Plant Biotechnology Research and Technology Transfer. Elsevier, Amsterdam.
3. **Han, R. & Ehlers, R.-U. (2000):** Pathogenicity, development, and reproduction of *Heterorhabditis bacteriophora* and *Steinernema carpocapsae* under axenic *in vivo* conditions. Journal of Invertebrate Pathology 75, 55-58.
4. **Holtmann, B., Kleine M. & Grundler, F.M.W. (2000):** Ultrastructure and anatomy of nematode induced syncytia in roots of susceptible and resistant sugar beet. Protoplasma 211, 39-50.
5. **Iraki, N., Salah, N., Sansour, M. A., Segal, D., Glazer, I., Johnigk, S.-A., Hussein, M.A. & Ehlers, R.-U. (2000):** Isolation and characterization of two entomopathogenic nematode strains, *Heterorhabditis indica* (Nematoda, Rhabditida), from the West Bank, Palestinian Territories. Journal of Applied Entomology 124, 375-380.
6. **Jeffke, T., Jende, D., Mätje, C., Ehlers, R.-U. & Berthe-Corti, L. (2000):** Growth of *Photorhabdus luminescens* in batch and glucose fed-batch culture. Applied Microbiology and Biotechnology 54, 326-330.
7. **Jessen, P., Strauch, O., Wyss, U., Luttmann, R. & Ehlers, R.-U. (2000):** Carbon dioxide triggers recovery from dauer juvenile stage in entomopathogenic nematodes (*Heterorhabditis* spp.) Nematology 2, 319-324.
8. **Lee, Y. K, Hippe-Sanwald, S., Lee, S.C., Hohenberg, H. & Hwang, B.K. (2000):** *In situ* localization of PR-1 mRNA and PR-1 protein in compatible and incompatible interactions of pepper stems with *Phytophthora capsici*. Protoplasma 211, 64-75.
9. **Lee, Y.K., Hong, J.K., Hippe-Sanwald, S. & Hwang, B.K. (2000):** Histological and ultrastructural comparisons of compatible, incompatible and DL- β -amino-n-butyric acid-induced resistance responses of pepper stems to *Phytophthora capsici*. Physiological and Molecular Plant Pathology 57, 269-280.
10. **Micha, S.G., Kistenmacher, S., Mölck, G. & Wyss, U. (2000):** Tritrophic interactions between cereals, aphids and parasitoids: Discrimination of different plant-host complexes by *Aphidius rhopalosiphi* (Hymenoptera: Aphidiidae). European Journal of Entomology 97, 539-543.

11. **Mölck, G., Pinn, H. & Wyss, U. (2000):** Manipulation of plant odour preference by learning in the aphid parasitoid *Aphelinus abdominalis* (Hymenoptera: Aphelinidae). *European Journal of Entomology* 97, 533-538.
12. **Petersen, G., Matthiesen, C., Francke, W. & Wyss, U. (2000):** Hyperparasitoid volatiles as possible foraging behaviour determinants in the aphid parasitoid *Aphidius uzbekistanicus* (Hymenoptera: Aphidiidae). *European Journal of Entomology* 97, 545-550.
13. **Puzio, P.S. & Grundler, F.M.W. (2000):** Isolation and application of nematode induced promoters, genes and proteins from *Arabidopsis thaliana*. In: G.E. de Vries, K. Metzlauff (eds.): *Phytosphere'99 - Highlights in European Plant Biotechnology Research and Technology Transfer*. Elsevier, Amsterdam.
14. **Puzio, P.S., Lausen, J., Heinen, P. & Grundler, F.M.W. (2000):** Promoter analysis of pyk 20, a gene from *Arabidopsis thaliana*. *Plant Science* 157, 245-255.
15. **Puzio, P.S., Neue, M., Grymaszewska, G., Ludwig-Müller, J. & Grundler, F.M.W. (2000):** *Plasmodiophora brassicae*-induced expression of pyk20, an *Arabidopsis* gene with glutamine-rich domain. *Physiological and Molecular Plant Pathology* 56, 79-84.
16. **Strauch, O. & Ehlers, R.-U. (2000):** Influence of the aeration rate on yields of the biocontrol nematodes *Heterorhabditis megidis* in monoxenic liquid cultures. *Applied Microbiology and Biotechnology* 54, 9-13.
17. **Strauch, O., Niemann, I., Neumann, A., Schmidt, A.J., Peters, A. & Ehlers, R.-U. (2000):** Storage and formulation of the entomopathogenic nematodes *Heterorhabditis indica* and *H. bacteriophora*. *Biocontrol* 45, 483-500.
18. **Verreet, J.-A., Klink, H. & Hoffmann, G.M. (2000):** Regional monitoring for disease prediction and optimization of plant protection measures: The IPM Wheat Model. *Plant Disease* 84, 816-826.
19. **Vierheilig, H., Bennett, R., Kiddle, G., Kaldorf, M. & Ludwig-Müller, J. (2000):** Differences in glucosinolate patterns and arbuscular mycorrhizal status of glucosinolate-containing plant species. *New Phytologist* 146, 343-352.
20. **Vierheilig, H., Garcia-Garrido M.J., Wyss, U. & Piché, Y. (2000):** Systemic suppression of mycorrhizal colonization in barley roots already colonized by AM-fungi. *Soil Biology and Biochemistry* 32, 589-595.
21. **Vierheilig, H., Maier, W., Wyss, U., Samson, J., Strack, D. & Piché, Y. (2000):** Cyclohexenone derivative- and phosphate-levels in split-root systems and their role in the systemic suppression of mycorrhization in precolonized barley plants. *Journal of Plant Physiology* 157, 593-599.
22. **Wolf, P.F.J., Verreet, J.-A., Maier, R. & Köhler, R. (2000):** An integrated pest management model (IPM sugar beet model) for threshold-oriented control of *Cercospora beticola* on sugar beet, developed under conditions in Southern Germany. In: M.J.C. Asher, B. Holtschulte, M. Richard-Molard, F. Rosso, G. Steinrücken, R. Beckers (eds.): *Advances in sugar beet research*, 103-121. International Institute for Beet Research, Brussels.

1999

1. **Ehlers, R.-U. & Niemann, I. (1999):** Molecular identification of *Photorhabdus luminescens* strains by amplification of specific fragments of the 16S ribosomal DNA. *Systematic and Applied Microbiology* 21, 509-519.
2. **Han, R.C. & Ehlers, R.-U. (1999):** Trans-specific nematicidal activity of *Photorhabdus luminescens*. *Nematology* 1, 687-693.
3. **Hass, B., Griffin, C.T. & Downes, M.J. (1999):** Persistence of *Heterorhabditis* infective juveniles in soil: Comparison of extraction and infectivity measurements. *Journal of Nematology* 31, 508-516.

4. **Hedke, K. & Verreet, J.-A. (1999):** Efficacy of single fungicide treatments in winter wheat using recommended and reduced dosages. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz* 106, 98-108.
5. **Johnigk, S.-A. & Ehlers, R.-U. (1999):** Juvenile development and life cycle of *Heterorhabditis bacteriophora* and *H. indica* (Nematoda: Heterorhabditidae). *Nematology* 1, 251-260.
6. **Johnigk, S.-A. & Ehlers, R.-U. (1999):** Endotokia matricida in hermaphrodites of *Heterorhabditis* spp. and the effect of the food supply. *Nematology* 1, 717-726.
7. **Jung, C. & Wyss, U. (1999):** New approaches to control plant parasitic nematodes. *Applied Microbiology and Biotechnology* 51, 439-446.
8. **Mölck, G., Micha, S.G. & Wyss, U. (1999):** Attraction to odour of infested plants and learning behaviour in the aphid parasitoid *Aphelinus abdominalis* (Hymenoptera: Aphelinidae). *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz* 106, 557-567.
9. **Pinior, A., Wyss, U., Piché, Y. & Vierheilig, H. (1999):** Plants colonized by an AM-fungus regulate further root colonization by AM-fungi through altered root exudation. *Canadian Journal of Botany* 77, 1-7.
10. **Puzio, P.S., Lausen J., Almeida-Engler, J., Cai, D.G., Gheysen, G. & Grundler, F.M.W. (1999):** Isolation of a gene from *Arabidopsis thaliana* related to nematode feeding structures. *Gene* 239, 163-172.
11. **Rühm, R. & Aumann, J. (1999):** The effect of Zn²⁺ on host finding and reproduction of *Heterodera schachtii*. *Nematology* 1, 443-444.
12. **Vierheilig, H., Böckenhoff, A., Knoblauch, M., Juge, C., van Bel, A.J.E., Grundler, F.M.W., Piché, Y. & Wyss, U. (1999):** *In vivo* observations of the arbuscular mycorrhizal fungus *Glomus mosseae* in roots by confocal laser scanning microscopy. *Mycological Research* 103, 311-314.
13. **Vierheilig, H., Garcia-Garrido, M. J., Wyss, U. & Piché, Y. (1999):** Systemic suppression of mycorrhizal colonization in barley roots precolonized by AM-fungi. *Soil Biology and Biochemistry* 32, 589-595.
14. **Wyss, U. (1999):** *Xiphinema index*, maintenance and feeding in monoxenic cultures. In: K. Maramorosch, F. Mahmood (eds.): *Maintenance of Human, Animal, and Plant Pathogen Vectors*, 253-281. Science Publishers, Enfield, NH, USA.

1998

1. **Abu Hatab, M., Gaugler, R. & Ehlers, R.-U. (1998):** Influence of culture method on *Steinernema glaseri* lipids. *Journal of Parasitology* 84, 215-221.
2. **Aumann, J., Dietsche, E., Rutencrantz, S. & Ladehoff, H. (1998):** Physico-chemical properties of the female sex pheromone of *Heterodera schachtii* (Nematoda: Heteroderidae). *International Journal for Parasitology* 28, 1691-1694.
3. **Aumann, J., Ladehoff, H. & Rutencrantz, S. (1998):** Gas chromatographic characterization of the female sex pheromone of *Heterodera schachtii* (Nematoda: Heteroderidae). *Fundamental and Applied Nematology* 21, 119-122.
4. **Christiansen-Weniger, P., Micha, S. G. & Siri, N. (1998):** Orientation of insects in a multitrophic system: the role of fragrances. In: P. J. Frosch, J. D. Johansen, I. R. White (eds.): *Fragrances, Beneficial and Adverse Effects*, 35-42. Springer, Berlin, Heidelberg, New York.
5. **Ehlers, R.-U. (1998):** Entomopathogenic nematodes - Safe biocontrol agents for sustainable systems. *Phytoprotection* 79, 94-102.
6. **Ehlers, R.-U., Lunau, S., Krasomil-Osterfeld, K. & Osterfeld, K.-H. (1998):** Liquid culture of the entomopathogenic nematode-bacterium complex *Heterorhabditis megidis/Photorhabdus luminescens*. *Biocontrol* 43, 77-96.
7. **Ehlers, R.-U. & Niemann, I. (1998):** Molecular identification of *Photorhabdus luminescens* strains by amplification of specific fragments of the 16S ribosomal DNA. *Systematic and Applied Microbiology* 21, 509-519.

8. **Grundler, F. M. W., Sobczak, M. & Golinowski, W. (1998):** Formation of wall openings in root cells of *Arabidopsis thaliana* following infection by the plant-parasitic nematode *Heterodera schachtii*. *European Journal of Plant Pathology* 104, 545-551.
9. **Han, R.C. & Ehlers, R.-U. (1998):** Cultivation of axenic *Heterorhabditis* spp. dauer juveniles and their response to non-specific *Photorhabdus luminescens* food signals. *Nematologica* 44, 425-435.
10. **Hussey, R. S. & Grundler, F. M. W. (1998):** Nematode parasitism of plants. In: R.N. Perry, D.J. Wright (eds.): *The Physiology and Biochemistry of Free-living and Plant-parasitic Nematodes*, 213-243. CAB International Press, Wallingford.
11. **Marticke, K.H., Reisener, H.J., Fischer, R. & Hippe-Sanwald, S. (1998):** *In situ* detection of a fungal glycoprotein-elicitor in stem rust-infected susceptible and resistant wheat using immunogold electron microscopy. *European Journal of Cell Biology* 76, 265-273.
12. **Perry, R. N. & Aumann, J. (1998):** Behaviour and sensory responses. In: R. N. Perry, D. J. Wright (eds.): *The Physiology and Biochemistry of Free-living and Plant-parasitic Nematodes*, 75-102. CAB International, Wallingford.
13. **Puzio, P.S., Cai, D.G., Ohl, S., Wyss, U. & Grundler, F. M. W. (1998):** Isolation of regulatory DNA regions related to differentiation of nematode feeding structures in *Arabidopsis thaliana*. *Physiological and Molecular Plant Pathology* 53, 177-193.
14. **Schnieder, F., Koch, G., Jung, C. & Verreet, J.-A. (1998):** The application of molecular markers for genetic characterization of *Septoria tritici* populations. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz* 105, 452-461.
15. **Strauch, O. & Ehlers, R.-U. (1998):** Food signal production of *Photorhabdus luminescens* inducing the recovery of entomopathogenic nematodes *Heterorhabditis* spp. in liquid culture. *Applied Microbiology and Biotechnology* 50, 369-374.
16. **Vierheilig, H., Coughlan, A., Wyss, U. & Piché, Y. (1998):** Ink and vinegar, a simple staining technique for arbuscular-mycorrhizal fungi. *Applied and Environmental Microbiology* 64, 5004-5007.
17. **Vierheilig, H. & Piché, Y. (1998):** A modified procedure for staining arbuscular mycorrhizal fungi in roots. *Zeitschrift für Pflanzenernährung und Bodenkunde* 161, 601-602.
18. **Wolf, P.F.J., Kraft, R. & Verreet, J.-A. (1998):** Schadrelevanz von *Cercospora beticola* (Sacc.) in Zuckerrüben als Grundlage einer Verlustprognose. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz* 105, 462-474.
19. **Wolf, P.F.J., Weis, F.J., Verreet, J.-A., Maier, H. & Tischner, H. (1998):** IPS (Integriertes Pflanzenschutzsystem)-Modell Zuckerrübe - Entwicklungsschritte und Einführung in die Praxis. *Gesunde Pflanzen* 50, 264-272.

1997

1. **Aumann, J. (1997):** Monoxenic culture of *Xiphinema index* (Nematoda: Longidoridae) on *Ficus carica*. *Nematologia Mediterranea* 25, 209-211.
2. **Barthels, N., van der Lee, F.M., Klap, J., Goddijn, O.M.J., Karimi, M., Puzio, P., Grundler, F.M.W., Ohl, S.A., Lindsey, K., Robertson, L., Robertson, W.M., van Montagu, M., Gheysen, G. & Sijmons, P.C. (1997):** Regulatory sequences of arabidopsis drive reporter gene expression in nematode feeding structures. *Plant Cell* 9, 2119-2134.
3. **Boemare, N., Neilson, K. & Ehlers, R.-U. (1997):** Preface - *Xenorhabdus-Steinernema* and *Photorhabdus-Heterorhabditis*, entomopathogenic bacterium nematode symbioses. *Symbiosis* 22, 1-3.
4. **Boenisch, A., Petersen, G. & Wyss, U. (1997):** Influence of the hyperparasitoid *Dendrocerus carpenteri* on the reproduction of the grain aphid *Sitobion avenae*. *Ecological Entomology* 22, 1-6.

5. **Börner, H. (1997):** Pflanzenkrankheiten und Pflanzenschutz, 7. Auflage. Eugen Ulmer, Stuttgart.
6. **Börner, H. (1997):** Unkrautbekämpfung. In: E.R. Keller, H. Hanus, K.-U. Heyland (eds.): Handbuch des Pflanzenbaues. Band 1: Grundlagen der landwirtschaftlichen Pflanzenproduktion, 362-394. Eugen Ulmer, Stuttgart.
7. **Cai, D.G., Kleine, M., Kifle, S., Harloff, H.J., Sandal, N.N., Marcker, K.A., Klein-Lankhorst, R.M., Salentijn, E.M.J., Lange, W., Stiekema, W.J., Wyss, U., Grundler, F.M.W. & Jung, C. (1997):** Positional cloning of a gene for nematode resistance in sugar beet. *Science* 275, 832-834.
8. **Ehlers, R.-U., Wulff, A. & Peters, A. (1997):** Pathogenicity of axenic *Steinernema feltiae*, *Xenorhabdus bovienii*, and the bacto-helminthic complex to larvae of *Tipula oleracea* (Diptera) and *Galleria mellonella* (Lepidoptera). *Journal of Invertebrate Pathology* 63, 212-217.
9. **Fenoll, C., Grundler, F.M.W. & Ohl, S.A., eds. (1997):** Cellular and Molecular Aspects of Plant-Nematode Interactions. Kluwer, Dordrecht.
10. **Giese, H., Hippe-Sanwald, S., Somerville, S. & Weller, J. (1997):** *Erysiphe graminis*. In: G.C. Carroll, P. Tudzynski (eds.): The Mycota V, Part B, Plant Relationships, 55-77. Springer, Berlin.
11. **Grundler, F.M.W. & Böckenhoff, A. (1997):** Physiology of nematode feeding and feeding sites. In: C. Fenoll, F.M.W. Grundler, S.A. Ohl (eds.): Cellular and Molecular Aspects of Plant-Nematode Interactions, 107-119. Kluwer, Dordrecht.
12. **Grundler, F.M.W., Sobczak, M. & Lange, S. (1997):** Defence responses of *Arabidopsis thaliana* during invasion and feeding site induction by the plant-parasitic nematode *Heterodera glycines*. *Physiological and Molecular Plant Pathology* 50, 419-429.
13. **Panstruga, R., Hippe-Sanwald, S., Lee, Y.K., Lataster, M., Lipka, V., Fischer, R., Liao, Y.C., Hausler, R.E., Kreuzaler, F. & Hirsch, H.J. (1997):** Expression and chloroplast-targeting of active phosphoenolpyruvate synthetase from *Escherichia coli* in *Solanum tuberosum*. *Plant Science* 127, 191-205.
14. **Peters, A. & Ehlers, R.-U. (1997):** Encapsulation of the entomopathogenic nematode *Steinernema feltiae* in *Tipula oleracea*. *Journal of Invertebrate Pathology* 69, 218-222.
15. **Peters, A., Gouge, D.H., Ehlers, R.-U. & Hague, N.G.M. (1997):** Avoidance of encapsulation by *Heterorhabditis* spp. infecting larvae of *Tipula oleracea*. *Journal of Invertebrate Pathology* 70, 161-164.
16. **Sobczak, M., Golinowski, W. & Grundler, F.M.W. (1997):** Changes in the structure of *Arabidopsis thaliana* roots induced during development of males of the plant parasitic nematode *Heterodera schachtii*. *European Journal of Plant Pathology* 103, 113-124.
17. **Stackebrandt, E., Ehlers, R.-U. & Rainey, F.A. (1997):** *Xenorhabdus* and *Photorhabdus*. Are they sister genera or are their members phylogenetically intertwined? *Symbiosis* 22, 59-65.
18. **Wolf, P.F.J. & Verreet, J.-A. (1997):** Epidemiologische Entwicklung von *Cercospora beticola* (Sacch.) in Zuckerrüben. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz* 104, 545-556.
19. **Wyss, U. (1997):** Root parasitic nematodes: An overview. In: C. Fenoll, F.M.W. Grundler, S.A. Ohl (eds.): Cellular and Molecular Aspects of Plant-Nematode Interactions, 5-22. Kluwer, Dordrecht.

1. **Böckenhoff, A., Prior, D.A.M., Grundler, F.M.W. & Oparka, K.J. (1996):** Induction of phloem unloading in *Arabidopsis thaliana* roots by the parasitic nematode *Heterodera schachtii*. *Plant Physiology* 112, 1421-1427.
2. **Dunier, M., Masson, J.P., Hokkanen, H. & Ehlers, R.-U. (1996):** OECD Workshop on the introduction of non-endemic nematodes for biological control: Scientific and regulatory policy issues. *Biocontrol Science and Technology* 6, 291-293.
3. **Ehlers, R.-U. (1996):** Current and future use of nematodes in biocontrol: Practice and commercial aspects with regard to regulatory policy issues. *Biocontrol Science and Technology* 6, 303-316.
4. **Ehlers, R.-U. & Hokkanen, H.M.T. (1996):** Insect biocontrol with non-endemic entomopathogenic nematodes (*Steinernema* and *Heterorhabditis* spp): Conclusions and recommendations of a combined OECD and COST Workshop on Scientific and Regulatory Policy Issues. *Biocontrol Science and Technology* 6, 295-302.
5. **Golinowski, W., Grundler, F.M.W. & Sobczak, M. (1996):** Changes in the structure of *Arabidopsis thaliana* during female development of the plant-parasitic nematode *Heterodera schachtii*. *Protoplasma* 194, 103-116.
6. **Grundler, F.M.W. (1996):** Engineering resistance against plant-parasitic nematodes. *Field Crops Research* 45, 99-109.
7. **Micha, S.G. & Wyss, U. (1996):** Aphid alarm pheromone (E)- β -farnesene: a host finding kairomone for the aphid primary parasitoid *Aphidius uzbekistanicus* (Hymenoptera: Aphidiidae). *Chemoecology* 7, 132-139.
8. **Sulistyanto, D. & Ehlers, R.-U. (1996):** Efficacy of the entomopathogenic nematodes *Heterorhabditis megidis* and *Heterorhabditis bacteriophora* for the control of grubs (*Phyllopertha horticola* and *Aphodius contaminatus*) in golf course turf. *Biocontrol Science and Technology* 6, 247-250.

1995

1. **Börner, H. (1995):** Unkrautbekämpfung. Gustav Fischer, Jena.
2. **Eggert, F., Höller, C., Luszyk, D. & Ferstl, R. (1995):** Identification of MHC-associated urinary chemosignals in mice. *Chemical Senses* 20, 124-125.
3. **Gerritsen, L.J.M., Vanderwolf, J.M., Vanvuurde, J.W.L., Ehlers, R.-U., Krasomil-Osterfeld, K.C. & Smits, P.H. (1995):** Polyclonal antisera to distinguish strains and form variants of *Photorhabdus (Xenorhabdus) luminescens*. *Applied and Environmental Microbiology* 61, 284-289.
4. **Hippe-Sanwald, S. (1995):** Low-temperature techniques as a tool in plant pathology. *Scanning Microscopy* 9, 881-899.
5. **Micha, S.G. & Wyss, U. (1995):** Bedeutung von Pflanzenduftstoffen bei der Wirtssuche von *Aphidius uzbekistanicus* (Hymenoptera, Aphidiidae), eines Parasitoiden der Großen Getreideblattlaus (*Sitobion avenae*). *Gesunde Pflanzen* 47, 300-307.
6. **Nobre, M.J.G., von Mende, N., Dolan, L., Schmidt, K.P., Evans, K. & Mulligan, B. (1995):** Immunolabeling of cell surfaces of *Arabidopsis thaliana* roots following infection by *Meloidogyne incognita* (Nematoda). *Journal of Experimental Botany* 46, 1711-1720.
7. **Rainey, F.A., Ehlers, R.-U. & Stackebrandt, E. (1995):** Inability of the polyphasic approach to systematics to determine the relatedness of the genera *Xenorhabdus* and *Photorhabdus*. *International Journal of Systematic Bacteriology* 45, 379-381.
8. **Wolf, P.F.J., Weis, F.J. & Verreet, J.-A. (1995):** Grundlagen einer integrierten Bekämpfung von *Cercospora beticola* in Zuckerrüben. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz* 102, 574-585.

1994

1. **Aumann, J. (1994):** The chemical nature of the amphidial and excretory system secretions of *Heterodera schachtii* (Nematoda, Heteroderidae) males. *Fundamental and Applied Nematology* 17, 186-189.
2. **Aumann, J. (1994):** Differential staining of "excretory" system mucins of *Heterodera schachtii* J₂-juveniles. *Nematologia Mediterranea* 22, 269-270.
3. **Böckenhoff, A. & Grundler, F.M.W. (1994):** Studies on the nutrient-uptake by the beet cyst-nematode *Heterodera schachtii* by *in-situ* microinjection of fluorescent probes into the feeding structures in *Arabidopsis thaliana*. *Parasitology* 109, 249-255.
4. **Börner, H., ed. (1994):** Pesticides in Ground and Surface Water. *Chemistry of Plant Protection*, Vol. 9. Springer, Berlin, Heidelberg, New York, Tokyo.
5. **Christiansen-Weniger, P. (1994):** Studies on semiochemicals affecting the host acceptance behaviour of *Asaphes vulgaris* Wlk. (Hymenoptera: Pteromalidae). *Norwegian Journal of Agricultural Sciences Suppl.* No. 16, 277-282.
6. **Clemens, C.D., Aumann, J., Spiegel, Y. & Wyss, U. (1994):** Attractant-mediated behaviour of mobile stages of *Heterodera schachtii*. *Fundamental and Applied Nematology* 17, 569-574.
7. **Grundler, F.M.W., Böckenhoff, A., Schmidt, K.-P., Sobczak, M., Golinowski, W. & Wyss, U. (1994):** *Arabidopsis thaliana* and *Heterodera schachtii*: A versatile model to characterize the interaction between host plants and cyst nematodes. In: F. Lamberti, C. de Giorgi, D.M. Bird (eds.): *Advances in Molecular Plant Nematology*, NATO AS1 Series A, Vol. 268, 171-180. Plenum, New York.
8. **Grundler, F.M.W. & Wyss, U. (1994):** Strategies of root parasitism by sedentary plant parasitic nematodes. In: K. Kohmoto, U.S. Singh, R.P. Singh (eds.): *Pathogenesis and Host Specificity in Plant Diseases*, Vol. II, Eukaryotes, 309-319. Pergamon Press, Oxford, 407 S.
9. **Grundler, F.M.W., Wyss, U. & Golinowski, W. (1994):** Sedentary endoparasitic nematodes in *Arabidopsis thaliana*. In: J. Bowman (ed.): *Arabidopsis: An Atlas of Morphology and Development*, 418-423. Springer, New York, 450 pp.
10. **Hardie, J., Hick, A.J., Höller, C., Mann, J., Merritt, L., Nottingham, S.F., Powell, W., Wadhams, L.J., Witthinrich, J. & Wright, A.F. (1994):** The responses of *Praon* spp. parasitoids to aphid sex-pheromone components in the field. *Entomologia Experimentalis et Applicata* 71, 95-99.
11. **Höller, C., Bargaen, H., Vinson, S.B. & Witt, D. (1994):** Evidence for the external use of juvenile hormone for host marking and regulation in a parasitic wasp, *Dendrocerus carpenteri*. *Journal of Insect Physiology* 40, 317-322.
12. **Höller, C., Micha, S.G., Schulz, S., Francke, W. & Pickett, J.A. (1994):** Enemy-induced dispersal in a parasitic wasp. *Experientia* 50, 182-185.
13. **Peters, A. & Ehlers, R.-U. (1994):** Susceptibility of leatherjackets (*Tipula paludosa* and *Tipula oleracea*, Tipulidae, Nematocera) to the entomopathogenic nematode *Steinernema feltiae*. *Journal of Invertebrate Pathology* 63, 163-171.
14. **Sijmons, P.C., Atkinson, H.J. & Wyss, U. (1994):** Parasitic strategies of root nematodes and associated host-cell responses. *Annual Review of Phytopathology* 32, 235-259.
15. **Strauch, O., Stoessel, S. & Ehlers, R.-U. (1994):** Culture conditions define automictic or amphimictic reproduction in entomopathogenic rhabditid nematodes of the genus *Heterorhabditis*. *Fundamental and Applied Nematology* 17, 575-582.
16. **Wolf, P.F.J. (1994):** Epidemiologisch begründetes Entscheidungsschema für eine integrierte Bekämpfung von *Drechslera tritici-repentis* (Died.) Shoem. (teleomorph *Pyrenophora tritici-repentis* (Died.) Drechs.). *Gesunde Pflanzen* 46, 164-172.

1993

1. **Aumann, J. (1993):** Chemosensory physiology of nematodes. *Fundamental and Applied Nematology* 16, 193-198.
2. **Aumann, J. (1993):** Permeability of chemosensillum-associated exudates for lectins in a plant-parasitic and in a free-living nematode. *Fundamental and Applied Nematology* 16, 381-382.
3. **Aumann, J. & Hashem, M. (1993):** Studies on substances with sex-pheromone activity produced by *Heterodera schachtii* females. *Fundamental and Applied Nematology* 16, 43-46.
4. **Aumann, J. & Hashem, M. (1993):** Effect of the formamidine acaricide chlordimeform on mobility and chemoreception of *Heterodera schachtii* males (Nematoda, Heteroderidae). *Nematologica* 39, 75-80.
5. **Borgemeister, C., Poehling, H.-M., Dinter, A. & Höller, C. (1993):** Effects of insecticides on life-history parameters of the aphid parasitoid *Aphidius rhopalosiphii* [Hym., Aphidiidae]. *Entomophaga* 38, 245-255.
6. **Ehlers, R.-U. & Gerwien, A. (1993):** Selection of entomopathogenic nematodes (Steinernematidae and Heterorhabditidae, Nematoda) for the biological control of crane-fly larvae *Tipula paludosa* (Tipulidae, Diptera). *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz* 100, 343-353.
7. **Hanf, C.-H. & Verreet, J.-A. (1993):** Consequences of a total ban on fungicide application on agriculture and agribusiness. In: J. Michalek, C.-H. Hanf (eds.): *The Economic Consequences of a Drastic Reduction in Pesticide Use in the EU. Revised Papers of a Workshop Held in Tannenfelde (Schleswig-Holstein), November 13th - 14th, 1993*, 205-226. Wissenschaftsverlag Vauk, Kiel.
8. **Höller, C., Bargaen, H., Vinson, S.B. & Braune, H.J. (1993):** Sources of the marking pheromones used for host discrimination in the hyperparasitoid *Dendrocerus carpenteri*. *Journal of Insect Physiology* 39, 649-656.
9. **Höller, C., Borgemeister, C., Haardt, H. & Powell, W. (1993):** The relationship between primary parasitoids and hyperparasitoids of cereal aphids - an analysis of field data. *Journal of Animal Ecology* 62, 12-21.
10. **Höller, C. & Haardt, H. (1993):** Low field performance of an aphid parasitoid, *Aphelinus abdominalis*, efficient in the laboratory (Hym., Aphelinidae). *Entomophaga* 38, 115-124.
11. **Höller, C. & Hörmann, R. (1993):** Patch marking in the aphid hyperparasitoid, *Dendrocerus carpenteri* - the information contained in patch marks. *Oecologia* 94, 128-134.
12. **Lunau, S., Stoessel, S., Schmidt-Peisker, A.J. & Ehlers, R.-U. (1993):** Establishment of monoxenic inocula for scaling-up in-vitro cultures of the entomopathogenic nematodes *Steinernema* spp. and *Heterorhabditis* spp. *Nematologica* 39, 385-399.
13. **Micha, S.G., Stammel, J. & Höller, C. (1993):** 6-Methyl-5-heptene-2-one, a putative sex and spacing pheromone of the aphid hyperparasitoid, *Alloxysta victrix* (Hymenoptera, Alloxystidae). *European Journal of Entomology* 90, 439-442.
14. **Verreet, J.-A. (1993):** Fungizide als Werkzeug des Integrierten Pflanzenschutzes im Weizenanbau. *Gesunde Pflanzen* 45, 135-154.

1992

1. **Aumann, J. (1992):** Effects of colchicine on mobility and chemoreception of *Heterodera schachtii* (Nematoda, Heteroderidae) males. *International Journal for Parasitology* 22, 87-90.

2. **Börner, H. & Zunke, U. (1992):** Praktikum der Phytopathologie. Paul Parey, Berlin, Hamburg.
3. **Endo, B.Y. & Wyss, U. (1992):** Ultrastructure of cuticular exudations in parasitic juvenile *Heterodera schachtii*, as related to cuticle structure. Protoplasma 166, 67-77.
4. **Haardt, H. & Höller, C. (1992):** Differences in life-history traits between isofemale lines of the aphid parasitoid *Aphelinus abdominalis* (Hymenoptera, Aphelinidae). Bulletin of Entomological Research 82, 479-484.
5. **Scholz, D. & Höller, C. (1992):** Competition for hosts between two hyperparasitoids of aphids, *Dendrocerus laticeps* and *Dendrocerus carpenteri* (Hymenoptera, Megaspilidae) - the benefit of interspecific host discrimination. Journal of Insect Behavior 5, 289-300.
6. **Voss, B., Utermöhl, P. & Wyss, U. (1992):** Variation between strains of the nematophagous endoparasitic fungus *Catenaria anguillulae* Sorokin. 2. Attempts to achieve parasitism of *Heterodera schachtii* Schmidt in pot trials. Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz 99, 311-318.
7. **Wyss, U. (1992):** Observations on the feeding behaviour of *Heterodera schachtii* throughout development, including events during moulting. Fundamental and Applied Nematology 15, 75-89.
8. **Wyss, U. & Grundler, F.M.W. (1992):** Feeding behaviour of sedentary plant parasitic nematodes. Netherlands Journal of Plant Pathology 98, 165-173.
9. **Wyss, U. & Grundler, F.M.W. (1992):** *Heterodera schachtii* and *Arabidopsis thaliana*, a model host-parasite interaction. Nematologica 38, 488-493.
10. **Wyss, U., Grundler, F.M.W. & Münch, A. (1992):** The parasitic behaviour of 2nd-stage juveniles of *Meloidogyne incognita* in roots of *Arabidopsis thaliana*. Nematologica 38, 98-111.
11. **Wyss, U., Voss, B. & Jansson, H.-B. (1992):** *In vitro* observations on the infection of *Meloidogyne incognita* eggs by the zoosporic fungus *Catenaria anguillulae* Sorokin. Fundamental and Applied Nematology 15, 133-139.

1991

1. **Aumann, J. (1991):** Effect of the sulfhydryl reagent, mersalyl acid, on sex-pheromone reception by *Heterodera schachtii* (Nematoda, Heteroderidae) males. Parasitology 102, 299-301.
2. **Aumann, J., Robertson, W.M. & Wyss, U. (1991):** Lectin binding to cuticle exudates of sedentary *Heterodera schachtii* (Nematoda: Heteroderidae) second stage juveniles. Revue de Nématologie 14, 113-118.
3. **Betka, M., Grundler, F. & Wyss, U. (1991):** Influence of changes in the nurse cell system (syncytium) on sex determination and development of the cyst nematode *Heterodera schachtii* - single amino-acids. Phytopathology 81, 75-79.
4. **Blossey, B. & Ehlers, R.-U. (1991):** Entomopathogenic nematodes (*Heterorhabditis* spp. and *Steinernema anomali*) as potential antagonists of the biological weed-control agent *Hyllobius transversovittatus* (Coleoptera, Curculionidae). Journal of Invertebrate Pathology 58, 453-454.
5. **Dijksterhuis, J., Harder, W., Wyss, U. & Veenhuis, M. (1991):** Colonization and digestion of nematodes by the endoparasitic nematophagous fungus *Drechmeria coniospora*. Mycological Research 95, 873-878.
6. **Ehlers, R.-U., Deseo, K.V. & Stackebrandt, E. (1991):** Identification of *Steinernema* spp. (Nematoda) and their symbiotic bacteria *Xenorhabdus* spp. from Italian and German soils. Nematologica 37, 360-364.
7. **Grundler, F., Betka, M. & Wyss, U. (1991):** Influence of changes in the nurse cell system (syncytium) on sex determination and development of the cyst nematode

Heterodera schachtii - total amounts of proteins and amino-acids. *Phytopathology* 81, 70-74.

8. **Grundler, F., Schnibbe, L. & Wyss, U. (1991):** *In vitro* studies on the behaviour of 2nd-stage juveniles of *Heterodera schachtii* (Nematoda, Heteroderidae) in response to host plant-root exudates. *Parasitology* 103, 149-155.
9. **Höller, C. (1991):** Evidence for the existence of a species closely related to the cereal aphid parasitoid *Aphidius rhopalosiphii* Destefani-Perez based on host ranges, morphological characters, isoelectric-focusing banding-patterns, cross-breeding experiments and sex-pheromone specificities (Hymenoptera, Braconidae, Aphidiinae). *Systematic Entomology* 16, 15-28.
10. **Höller, C., Williams, J.J. & Vinson, S.B. (1991):** Evidence for a two-component external marking pheromone system in an aphid hyperparasitoid. *Journal of Chemical Ecology* 17, 1021-1035.
11. **Nissen, U., Laborius, G.-A. & Schulz, F.-A. (1991):** Populationsvergleich von *Prostephanus truncatus* (Horn) (Col., Bostrichidae) unterschiedlicher geographischer Herkunft. *Journal of Applied Entomology* 112, 124-137.
12. **Reuter, G., Aumann, J., Wyss, U., Jansson, H.-B. & Schauer, R. (1991):** *Panagrellus redivivus* - Failure to find evidence for the occurrence and biosynthesis of sialic acids. *Experimental Parasitology* 73, 389-395.
13. **Sijmons, P.C., Grundler, F.M.W., von Mende, N., Burrows, P.R. & Wyss, U. (1991):** *Arabidopsis thaliana* as a new model host for plant-parasitic nematodes. *Plant Journal* 1, 245-254.
14. **Trudgill, D.L., Robertson, W.M. & Wyss, U. (1991):** Analysis of the feeding of *Xiphinema diversicaudatum*. *Revue de Nématologie* 14, 107-112.

1990

1. **Aumann, J., Clemens, C.D. & Wyss, U. (1990):** Influence of lectins on female sex pheromone reception by *Heterodera schachtii* (Nematoda: Heteroderidae) males. *Journal of Chemical Ecology* 16, 2371-2380.
2. **Basedow, T., Schulz, F.-A. & Al-Najjar, A. (1990):** Die Auswirkungen abgestufter Produktionsintensitäten auf Schaderreger, Erträge und Wirtschaftlichkeit. II. Ökonomische Aspekte. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz* 97, 284-298.
3. **Börner, H. (1990):** Veränderungen im Ergosterolgehalt bei *Sclerotinia sclerotiorum* während der Sklerotienentwicklung und karpogenen Keimung. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz* 97, 306-312.
4. **Börner, H. (1990):** *Pflanzenkrankheiten und Pflanzenschutz*, 6. Auflage. Eugen Ulmer, Stuttgart.
5. **Ehlers, R.-U., Stoessel, S. & Wyss, U. (1990):** The influence of phase variants of *Xenorhabdus* spp. and *Escherichia coli* (Enterobacteriae) on the propagation of entomopathogenic nematodes of the genera *Steinernema* and *Heterorhabditis*. *Revue de Nématologie* 13, 417-424.
6. **Höller, C. (1990):** Overwintering and hymenopterous parasitism in autumn of the cereal aphid *Sitobion avenae* (F.) in Northern FR Germany. *Zeitschrift für Angewandte Entomologie* 109, 21-28.
7. **Pütz, J., Meinert, F., Wyss, U., Ehlers, R.-U. & Stackebrandt, E. (1990):** Development and application of oligonucleotide probes for molecular identification of *Xenorhabdus* species. *Applied and Environmental Microbiology* 56, 181-186.
8. **Sanft, U. & Wyss, U. (1990):** Monoxenic test for the resistance of potato genotypes to the potato cyst nematode *Globodera pallida*. 1. Cultivation of the genotypes. *Potato Research* 33, 43-53.

9. **Sanft, U. & Wyss, U. (1990):** Monoxenic test for the resistance of potato genotypes to the potato cyst nematode *Globodera pallida*. 2. Performance of the test with nematodes. Potato Research 33, 55-66.
10. **Sanft, U. & Wyss, U. (1990):** Monoxenic test for the resistance of potato genotypes to the potato cyst nematode *Globodera pallida*. 3. Use and examination of the test. Potato Research 33, 67-76.
11. **Voss, B. & Wyss, U. (1990):** Variation between strains of the nematophagous endoparasitic fungus *Catenaria anguillulae* Sorokin. 1. Factors affecting parasitism *in vitro*. Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz 97, 416-430.
12. **Zunke, U. (1990):** Ectoparasitic feeding behaviour of the root lesion nematode, *Pratylenchus penetrans*, on root hairs of different host plants. Revue de Nématologie 13, 331-337.

1989

1. **Al-Najjar, A., Basedow, T. & Schulz, F.-A. (1989):** Effects of gradually differing intensities of growing wheat on pests and diseases, yields and economy. 1. Phytomedical aspects. Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz 96, 561-584.
2. **Aumann, J. (1989):** Enzymatic effects on lectin binding to *Heterodera schachtii* (Nematoda, Heteroderidae) males. Nematologica 35, 461-468.
3. **Aumann, J. & Wyss, U. (1989):** Histochemical studies on exudates of *Heterodera schachtii* (Nematoda: Heteroderidae) males. Revue de Nématologie 12, 309-315.
4. **Finck, G. (1989):** Der Einsatz von Umwandlungsprodukten des Kalkstickstoffs im Freiland zur Verhinderung der Sklerotienkeimung von *Sclerotinia sclerotiorum*, dem Erreger der Weißstengeligkeit. Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz 96, 508-520.
5. **Perry, R.N., Zunke, U. & Wyss, U. (1989):** Observations on the response of the dorsal and subventral oesophageal glands of *Globodera rostochiensis* to harching stimulation. Revue de Nématologie 12, 91-96.
6. **Purrini, K. & Keil, H. (1989):** *Ascogregarina bostrichidorum* n. sp. (Lecudinidae, Eugregarinida), a new gregarine parasitizing the larger grain borer, *Prostephanus truncatus* Horn (1878) (Bostrichidae, Coleoptera). Archiv für Protistenkunde 137, 165-171.
7. **Veenhuis, M., Vanwijk, C., Wyss, U., Nordbring-Hertz, B. & Harder, W. (1989):** Significance of electron dense microbodies in trap cells of the nematophagous fungus *Arthrobotrys oligospora*. Antonie van Leeuwenhoek Journal of Microbiology 56, 251-261.

1988

1. **Ehlers, R.-U., Wyss, U. & Stackebrand, E. (1988):** 16S Ribosomal RNA cataloging and the phylogenetic position of the genus *Xenorhabdus*. Systematic and Applied Microbiology 10, 121-125.
2. **Höller, C. & Braune, J. (1988):** The use of isoelectric focusing to assess percentage hymenopterous parasitism in aphid populations. Entomologia Experimentalis et Applicata 47, 105-114.
3. **Leliveldt, B., Laborius, G.-A. & Schulz, F.-A. (1988):** The distribution of *Mattesia* sp. (Neogregarinida, Ophryocystidae), a pathogen of *Prostephanus truncatus* (Horn) (Coleoptera, Bostrichidae) in Togo. Journal of Stored Products Research 24, 241-243.
4. **Wyss, U. (1988):** Pathogenesis and host-parasite specificity in nematodes. In: R.S. Singh, U.S. Singh, W.M. Hess, D.J. Weber (eds.): Experimental and Conceptual Plant Pathology, Vol. 2, 417-432. Gordon and Breach, New York.

5. **Wyss, U., Robertson, W.M. & Trudgill, D.L. (1988):** Oesophageal bulb function of *Xiphinema index* and associated root cell responses, assessed by video-enhanced contrast light microscopy. *Revue de Nématologie* 11, 253-261.

1987

1. **Aumann, J. & Wyss, U. (1987):** Lectin binding sites on mobile stages of *Heterodera schachtii* Schmidt (Nematoda, Heteroderidae). *Nematologica* 33, 410-418.
2. **Basedow, T., Beckmann, C. & Runge, I. (1987):** Problematik von Freilandversuchen zur Prüfung der Wirkung von Pflanzenschutzmitteln auf epigäische Raubarthropoden im Ackerbau. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz* 94, 260-275.
3. **Börner, H. (1987):** Veränderungen der Polyamingehalte während der Sklerotienkeimung und Apothecienentwicklung von *Sclerotinia sclerotiorum*, dem Erreger des Rapskrebsees. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz* 94, 190-196.
4. **Wyss, U. (1987):** Video assessment of root cell responses to dorylaimid and tylenchid nematodes. In: J.A. Veech, D.W. Dickson (eds.): *Vistas on Nematology*, 211-220. Society of Nematologists, Hyattsville, MD.

1986

1. **Isenbeck, M. & Schulz, F.-A. (1986):** Biological control of fireblight (*Erwinia amylovora* [Burr] Winslow et al.) on ornamentals. 2. Investigation about the mode of action of the antagonistic bacteria. *Phytopathologische Zeitschrift* 116, 308-314.
2. **Schulz, F.-A. & Ceynowa, J. (1986):** Occurrence and toxicological activity of molds on cereals stored in airtight containers. *Phytoparasitica* 14, 333.
3. **Schulz, F.-A. & Laborius, G.-A. (1986):** Strategy for bio-integrated control of *Prostephanus truncatus*. *Phytoparasitica* 14, 341-342.
4. **Wyss, U. (1986):** Nurse cell systems of dorylaimid and tylenchid nematodes. *Journal of Nematology* 18, 598.
5. **Wyss, U. & Zunke, U. (1986):** The potential of high-resolution video-enhanced contrast microscopy in nematological research. *Revue de Nématologie* 9, 91-94.
6. **Wyss, U. & Zunke, U. (1986):** Gnotobiology of cyst nematodes, its potential in basic research. In: F. Lamberti, C.E. Taylor (eds.): *Cyst Nematodes*, 147-162. Plenum, New York.
7. **Wyss, U. & Zunke, U. (1986):** Observations on the behaviour of second-stage juveniles of *Heterodera schachtii* inside host roots. *Revue de Nématologie* 9, 153-165.
8. **Zunke, U., Rössner, J. & Wyss, U. (1986):** Feeding behaviour of *Aphelenchoides hamatus* on five different phytopathogenic fungi and *Agaricus campestris*. *Nematologica* 32, 194-201.

1985

1. **Börner, H. (1985):** Nachweis von Arginase in den Sklerotien und Apothecien von *Sclerotinia sclerotiorum*, dem Erreger der Weißstängeligkeit des Rapses. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz* 92, 204-208.
2. **Börner, H. (1985):** Berichtigung. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz* 92, 336.
3. **Finck, G. & Börner, H. (1985):** Beeinflussung der Sklerotienkeimung von *Sclerotinia sclerotiorum*, dem Erreger der Weißstängeligkeit des Rapses, durch

Umwandlungsprodukte von Kalkstickstoff. Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz 92, 449-454.

4. **Gleim, D. (1985):** Beeinflussung der mikrobiellen Aktivität eines lehmigen Sandes durch Pflanzenschutzmittel für den Wintergerstenanbau. Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz 92, 568-580.
5. **Isenbeck, M. & Schulz, F.-A. (1985):** Biological control of fireblight (*Erwinia amylovora* [Burr] Winslow et al.) on ornamentals. 1. Control of the pathogen by antagonistic bacteria. Phytopathologische Zeitschrift 113, 324-333.
6. **Plieth, H.-E. & Börner, H. (1985):** Der Abbau von Pendimethalin im Boden von Gerstenkulturen unter dem Einfluss anderer Pflanzenschutzmittel und von Stroh. Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz 92, 288-304.
7. **Riecken, I. (1985):** Versuche zur Verhütung der Rotpustelkrankheit (*Nectria cinnabarina*). Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz 92, 516-529.
8. **Zunke, U. (1985):** Zur Bildung der subkristallinen Schicht bei *Heterodera schachtii* unter aseptischen Bedingungen. Nematologica 31, 117-120.

1984

1. **Krauthausen, H.-J. & Wyss, U. (1984):** Microanalysis of proteins at feeding sites of the developing cyst nematode *Heterodera schachtii*. Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz 91, 411-423.
2. **Wyss, U., Stender, C. & Lehmann, H. (1984):** Ultrastructure of feeding sites of the cyst nematode *Heterodera schachtii* Schmidt in roots of susceptible and resistant *Raphanus sativus* var. *oleiformis* Pers. cultivars. Physiological Plant Pathology 25, 21-37.

1983

1. **Börner, H. (1983):** Pflanzenkrankheiten und Pflanzenschutz, 5. Auflage. Eugen Ulmer, Stuttgart.
2. **Wyss, U. (1983):** Recognition and specificity in nematode diseases. Phytopathology 73, 769.

1982

1. **Kabsch, U. (1982):** Veränderungen im Kohlenhydratstoffwechsel in mit *Sphaerotheca fuliginea* infizierten Gurkenblättern. Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz 89, 113-124.

1981

1. **Börner, H. (1981):** Pflanzenkrankheiten und Pflanzenschutz, 4. Auflage. Eugen Ulmer, Stuttgart.
2. **Börner, H. & Ajang, M.R. (1981):** Die Auswaschung von Paraquat aus den Böschungen von Entwässerungsgräben nach chemischen Grabenentkrautungsmaßnahmen. Nachrichtenblatt des Deutschen Pflanzenschutzdienstes 33, 55-59.
3. **Rudolph, H., Thönes, S. & Kabsch, U. (1981):** Analyse der löslichen Kohlenhydrate während der durch Kältebehandlung induzierten Synthese der Sphagnorubine. Zeitschrift für Pflanzenphysiologie 101, 323-331.

1979

1. **Börner, H. (1979):** Ursachen der selektiven Wirkung der Photosynthesehemmer Phenmedipham und Bentazon. Zeitschrift für Naturforschung C 34, 926-930.
2. **Koch, O. & Schulz, F.-A. (1979):** Isoenzymenspektren der Pektin-Depolymerase von *Glomerella cingulata* und *Pezizula malicorticis*. Phytopathologische Zeitschrift 94, 261-269.

1978

1. **Börner, H. (1978):** Pflanzenkrankheiten und Pflanzenschutz, 3. Auflage. Eugen Ulmer, Stuttgart.
2. **Schröder, C.M. (1978):** Serologische Methoden zum Nachweis des Erregers des Feuerbrandes, *Erwinia amylovora*. Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz 85, 393-403.
3. **Schröder, C.M. (1978):** Untersuchungen zur Epidemiologie des Feuerbrandes, verursacht durch *Erwinia amylovora*. Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz 85, 482-501.
2. **Schulz, F.-A. (1978):** Some physiological and biochemical aspects of action mechanism of fungal parasites during fruit storage. Fruits 33, 15-21.

1977

1. **Börner, H. (1977):** Untersuchungen über das Verhalten von Paraquat nach Anwendung bei der chemischen Grabenentkrautung. Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz 84, 705-718.
2. **Linß, H. (1977):** Untersuchungen über die Veränderungen des Stoffwechsels der Rapspflanzen nach Infektion mit *Plasmodiophora brassicae* Wor. Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz 84, 497-508.

1975

1. **Börner, H. (1975):** Pflanzenkrankheiten und Pflanzenschutz, 2. Auflage. Eugen Ulmer, Stuttgart.

1974

1. **Koch, O. & Schulz, F.-A. (1974):** Einfluß ökologischer Parameter auf die Bildung multipler Formen von Pektinasen durch *Pezizula malicorticis* in vitro. Phytopathologische Zeitschrift 81, 184-187.

1971

1. **Börner, H. (1971):** Pflanzenkrankheiten und Pflanzenschutz. Eugen Ulmer, Stuttgart.
2. **Menck, B.H. & Börner, H. (1971):** Die Verbreitung des Ackerfuchsschwanzes (*Alopecurus myosuroides* Huds.) in Schleswig-Holstein. Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz 78, 217-228.
3. **Niehuss, M. & Börner, H. (1971):** Untersuchungen über den Einfluß von Herbiziden auf Fische. Nachrichtenblatt des Deutschen Pflanzenschutzdienstes 23, 113-117.
4. **Niehuss, M. & Börner, H. (1971):** Persistenz und Verteilung von Diuron, Dichlobenil, Dalapon und Paraquat nach Einsatz in Entwässerungsgräben. Ein Beitrag zum Problem der chemischen Grabenentkrautung. Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz 78, 385-397.

5. **Prante, G. & Börner, H. (1971):** Die Verbreitung des Flughafers (*Avena fatua* L.) in Schleswig-Holstein. Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz 78, 416-428.

1969

1. **Börner, H., Burgemeister, H. & Schroeder, M. (1969):** Untersuchungen über Aufnahme, Verteilung und Abbau von Harnstoffherbiziden durch Kulturpflanzen, Unkräuter und Mikroorganismen. Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz 76, 385-395.